



THE JUNGLE CROW
(*Corvus coronoides levallanti*)



CSL

BIRDS OF AN INDIAN GARDEN

BY

T. BAINBRIGGE FLETCHER, R.N., F.L.S., F.E.S., F.Z.S.

Imperial Entomologist

AND

C. M. INGLIS, M.B.O.U., F.E.S., F.Z.S.

Curator, Natural History Museum, Darjiling

With 30 Coloured Plates and numerous Illustrations in the Text

CALCUTTA & SIMLA
THACKER, SPINK & CO.

1924

2601
19.11.25



CSL

012475

598.2954
F57B

PUBLISHERS' NOTE

The coloured plates and text of these articles originally appeared in the *Agricultural Journal of India* during the years 1919-1924 under the title of "SOME COMMON INDIAN BIRDS."

It has been found necessary to alter this title to "BIRDS OF AN INDIAN GARDEN" to avoid confusion with works of similar titles by other authors.

The copyright is vested in the Secretary of State for India by whom all rights of reproduction and translation are reserved.

12470



PREFACE.

THIS series of papers dealing with some of the more common of our Indian Birds first appeared in the *Agricultural Journal of India* during the years 1919-1924 and had its inception in the desire to help the recognition of the more important birds, from the agricultural view-point, by the issue of a series of coloured plates. There was considerable difficulty in getting this done as work of this nature requires an artist who is also an ornithologist; otherwise, the work produced is likely to be untrue to nature. This combination of qualities was unattainable for some time and the scheme had perforce to be laid aside for some years until Mr. C. M. Inglis, a well-known worker on Indian Ornithology for many years past, visited Pusa and showed me some of his sketches of birds and insects. These indicated that Mr. Inglis was in a position to provide us with the coloured plates required and, on proposing this to him, he readily fell in with the idea. The main share of such popularity as these papers have achieved during their progress in the *Agricultural Journal of India* has been due, I am sure, to the excellent series of plates printed from Mr. Inglis' original paintings. My share has been to provide some appropriate letterpress and, in this reissue of the papers, to provide the uncoloured illustrations, most of which have been prepared by my artists directly from specimens or have been copied from original photographs or, in a few cases, from previous books on Indian Birds.

In the original issue we were forced to publish the articles without any regard to systematic order, but just as the plates happened to be completed and in explanation of this I wrote at the time.

"The haphazard manner in which we are dealing with our common birds may appear to demand some sort of apology, but the apparent lack of arrangement is simply due to nothing but the relative



rate at which it is possible to give suitable illustrations of the various birds dealt with. Generally speaking, it is necessary to paint the plate of each feathered friend or foe, with its natural leafy or other background, including in many cases its nest and eggs, at a particular time of the year, and it may chance for various reasons that it is not possible to secure a life-like portrait of any particular bird in its natural haunts in time to enable a plate to be painted just when it is required." The observant will doubtless see more in this than meets the casual eye.

In reissuing the series, however, the opportunity has been taken to rearrange the papers in systematic order and to amend the nomenclature in accordance with Mr. Stuart-Baker's Hand-list of Indian Birds, recently published in the *Bombay Natural History Society's Journal* and which epitomizes the revised issue of the volumes on Birds now in course of publication in the *Fauna of British India* series.

Although numerous illustrations, coloured and otherwise, of Indian Birds have appeared from time to time in many books and articles, both scientific and popular, such illustrations have been almost wholly devoted to figuring new and rare species, in the case of the more purely ornithological works, or species of particular groups, such as Mr. Stuart-Baker's excellent series of coloured plates illustrating the Indian Ducks and Pigeons, or, in the case of professedly popular publications which have included figures of the commoner birds, such illustrations have been rather crude and unsatisfactory—often, indeed, barely recognizable. The present publication will therefore, I hope, fill a gap by enabling the non-ornithological reader, who knows little or nothing about Indian Birds, to recognize and hence to take an interest in those commoner species which are to be met with in practically all parts of the country. There must be many in India, both Indians and Europeans, residents and visitors, who derive deep, albeit simple, joy from watching the habits of the wild bird-life which can be seen at any season of the year in every compound in the country, but who have



often felt themselves debarred from such pleasures by their want of knowledge of the birds concerned—their names and what is already known about them. For such readers, in particular, this little book is intended. In Western countries, where every kind of bird is well-known and where cheap and well-illustrated books on Birds are readily obtainable, observers of wild bird-life are numbered by the hundred and an immense mass of information has been accumulated on the occurrence and habits of all the native birds. In India, however, except for the very small number who may be dubbed ornithologists, such observers are themselves *rare aves*; yet India offers an immense field for observation of this sort. I do not refer to the mere collecting of skins and nests and eggs; nor must the epithet “mere” be taken to imply disparagement of scientific advance in these directions. But there is no locality in India in which very useful observations cannot be made on even our commonest kinds of birds. Take the following points for instance. Many of our birds are local migrants: on what dates in each locality do they appear and leave again? In what nests do the local cuckoos deposit their eggs and how do they accomplish this? What is the normal food of our common birds and what natural causes keep them in check? Such questions, and many others, can only be answered by local observers, not in one place or in one year, but in many localities throughout the country. All accurate observations, then, on bird-life will be valuable in proportion to their number, provided of course that they are accessible. This implies publication for which there is no lack of facilities in India: it is only necessary to quote, as one example, the *Journal of the Bombay Natural History Society*. Every number of this Journal contains articles and short notes on the occurrence and habits of Indian Birds and everyone in India who is interested in bird-life is strongly recommended to join the Society and to read the Journal and contribute their own notes to it.

There is an old English saw anent the difficulty of getting a quart into a pint pot and the same principle necessarily applies to a



book of this nature. About two hundred species of Indian Birds might be described as of common occurrence in the Plains of India and of these a large proportion may be seen, at one time or another, in an average Indian Garden: it was out of the question to include all of these and therefore only "some" of the common birds could be selected for inclusion. It is hoped, however, that this selection will serve as an introduction to the study of Indian Birds. To those desiring a concise handbook to the commoner species I would recommend Mr. Dewar's "Indian Birds: a key to the common birds of the Plains of India" and for those whose keenness leads them still further there are the volumes on Birds in the *Fauna of British India* series, of which a revised edition is now in course of issue.

T. BAINBRIGGE FLETCHER.

PUSA:
March 1924.

"..... Over the spangled grass
Swept the swift footsteps of the lovely Light,
Turning the tears of Night to joyous gems,
Decking the earth with radiance, 'brodering
The sinking storm-clouds with a golden fringe,
Gilding the feathers of the palms, which waved
Glad salutation; darting beams of gold
Into the glades; touching with magic wand
The stream to rippled ruby; in the brake
Finding the mild eyes of the antelopes
And saying 'It is day'; in nested sleep
Touching the small heads under many a wing
And whispering 'Children, praise the light of day.'
Whereat there piped anthems of all the birds,
The Koel's fluted song, the Bulbul's hymn,
The 'morning, morning' of the painted thrush,
The twitter of the sunbirds starting forth
To find the honey ere the bees be out,
The grey crow's caw, the parrot's scream, the strokes
Of the green hammersmith, the myna's chirp,
The never-finished love-talk of the doves."

Light of Asia.

CONTENTS

List of Articles and Coloured Plates.

Plate No. I	JUNGLE CROW (<i>Corvus coronoides levaillanti</i>)	Frontispiece
		Facing Page
" " II	INDIAN HOUSE-CROW (<i>Corvus splendens</i>) ..	7
" " III	BENGAL TREE-PIE (<i>Dendrocitta rufa vagabunda</i>) ..	15
" " IV	BENGAL JUNGLE-BABBLER OR "SEVEN SISTERS" (<i>Turdoides terricolor terricolor</i>) ..	21
" " V	BENGAL RED-VENTED BULBUL (<i>Molpastes</i> <i>haemorrhous bengalensis</i>) ..	27
" " VI	BLACK DRONGO OR KING CROW (<i>Dicrurus</i> <i>macrocerus</i>) ..	31
" " VII	TAILOR-BIRD (<i>Orthotomus sutorius sutorius</i>) ..	37
" " VIII	BLACK-HEADED ORIOLE (<i>Oriolus luteolus luteolus</i>) ..	45
" " IX	COMMON MYNA (<i>Acridotheres tristis tristis</i>) ..	49
" " X	PIED MYNA (<i>Sturnopastor contra contra</i>) ..	55
" " XI	INDIAN PARADISE FLY-CATCHER (<i>Terpsiphone</i> <i>paradisi paradisi</i>) ..	59
" " XII	INDIAN MAGPIE-ROBIN OR DAYAL (<i>Copsychus</i> <i>saularis saularis</i>) ..	65
" " XIII	WEAVER-BIRD OR BAYA (<i>Ploceus philippinus</i>) ..	69
" " XIV	INDIAN HOUSE-SPARROW (<i>Passer domesticus indicus</i>) ..	75
" " XV	WHITE WAGTAIL (<i>Motacilla alba</i>) ..	79
" " XVI	INDIAN TREE-PIBIT (<i>Anthus hodgsoni</i>) ..	83
" " XVII	INDIAN WHITE-EYE (<i>Zosterops palpebrosa palpebrosa</i>) ..	87
" " XVIII	PURPLE SUN-BIRD OR HONEY-SUCKER (<i>Cyrtostomus asiaticus asiaticus</i>) ..	91
" " XIX	NORTHERN GOLDEN-BACKED WOODPECKER (<i>Brachypternus aurantius aurantius</i>) ..	99
" " XX	NORTHERN GREEN BARBET (<i>Thereicerya</i> <i>zeylanicus caniceps</i>) ..	103
" " XXI	COPPERSMITH (<i>Xantholæma hæmacephala indica</i>) ..	107
" " XXII	NORTHERN INDIAN ROLLER (<i>Coracias benghalensis</i> <i>benghalensis</i>) ..	111
" " XXIII	COMMON INDIAN BEE-EATER (<i>Merops orientalis</i> <i>orientalis</i>) ..	117
" " XXIV	BLUE-TAILED BEE-EATER (<i>Merops superciliosus</i> <i>javanicus</i>) ..	123
" " XXV	INDIAN HOOPOE (<i>Upupa epops orientalis</i>) ..	127
" " XXVI	INDIAN LONG-TAILED NIGHTJAR (<i>Caprimulgus</i> <i>macrurus albonotatus</i>) ..	137
" " XXVII	HAWK-CUCKOO (<i>Hierococcyx varius</i>) ..	143
" " XXVIII	ROSE-RINGED PAROQUET (<i>Psittacula torquata</i>) ..	149
" " XXIX	INDIAN SPOTTED OWLET (<i>Carine noctua brama</i>) ..	155
" " XXX	CATTLE EGRET (<i>Bubulcus ibis coromandus</i>) ..	159

*List of Illustrations in the Text.*

	PAGE
JUNGLE CROW	3
A KASHMIRI SONGSTER	6
HOUSE-CROW	9
TREE-PIE	17
INDIAN MAGPIE	20
"SEVEN SISTERS"	23
COMMON KINGFISHER	26
DRONGO	33
NEST OF THE TAILOR-BIRD	40
A BEAUTIFULLY CONSTRUCTED NEST	44
COMMON KINGFISHER	54
A YOUNG BABBLER	58
RED-WHISKERED BULBUL	64
FEATHERS OF MAGPIE-ROBIN	66
COMMON MYNA	68
FEATHERS FROM TAIL AND WING OF FEMALE KOËL	74
WHITE WAGTAIL	80
YOUNG BABBLER	82
HEN KOËL	86
THE DAYAL	90
THE BLACK-HEADED ORIOLE	98
GOLDEN-BACKED WOODPECKER	101
THE BLUE-THROATED BARBET	106
BRAIN-FEVER BIRD	110
INDIAN ROLLER	112
THE KING CROW	116
COMMON BEE-EATER	118
A FEMALE HONEY-SUCKER	122
COMMON BENGAL BULBUL	126
HEAD OF EUROPEAN HOOPOE	128
COMMON INDIAN NIGHTJAR, LEFT FOOT AND HEAD	139
ROSE-RINGED PAROQUET	142
HEAD AND FOOT OF HAWK-CUCKOO	144
SPOTTED DOVE	148
ROSE-RINGED PAROQUETS ATTACKING WILD FRUITS	150
SUGARCANE DAMAGED BY PARROTS IN ASSAM	151
HEAD OF MALE ROSE-RINGED PAROQUET	154
CATTLE EGRET, <i>winter plumage</i>	160
CATTLE EGRET, <i>summer plumage</i>	161
BIRD'S EGGS, <i>Plate I</i>	162
" " " <i>II</i>	163
	<i>Facing page</i>



THE JUNGLE CROW

(*Corvus coronoides levaillanti*).

As I was walking all my lane
I heard twa corbies makin' a mane,
The tain until the tither say,
"Whar'll we gang and dine the day?"

Border Ballad.

"This is the end that comes
To all," quoth Channa; "he upon the pyre
Whose remnants are so petty that the crows
Caw hungrily, then quit the fruitless feast."

The Light of Asia.

THE "country cousin" of the House Crow is the Jungle Crow which, although it seems to prefer a *mofussil* life, is often seen, like human country-folk, in large towns to which it comes on a shorter or longer visit or in which it is frequently a permanent dweller. The House Crow and the Jungle Crow are the only two crows found commonly distributed in the Plains of India and are easily distinguishable, the House Crow being a smaller bird with a grey neck, the Jungle Crow being larger, with a heavier bill and black neck. In the extreme North, in areas bounding the Hills, other species of *Corvus*, such as the Raven, the Carrion-Crow, the Eastern Rook and the Eastern Jackdaw, descend to the Plains in the winter months. Of these, the Raven, distinguished by its much larger size, is common throughout the year as far South as the United Provinces; the Carrion-Crow, found along the North-West Frontier, is much like the Jungle Crow but with more intensely black plumage; the Eastern Rook, which is also somewhat like the Jungle Crow but has a more slender bill, the base of which is bare



of feathers in adult birds, and the lower plumage intensely glossed with blue and purple, is also found along the North-West Frontier and often does considerable good by feeding on grasshoppers and other insects on cultivated land; whilst the Eastern Jackdaw, which looks more like the House Crow but has the chin and throat grey like the breast, is found in the winter as far South as Ferozepur and as far East as Ambala.

The Jungle Crow is found throughout the Plains of India, and has been divided into forms, distinguished by length of wing and bill, the medium-sized form being the subject of our Plate, whilst in Southern India and Ceylon it is replaced by the Southern Jungle Crow (*Corvus coronoides culminatus*) with much shorter wing and smaller, slenderer bill; in the Himalaya it is replaced by the Himalayan Jungle Crow (*Corvus coronoides intermedius*), a larger race with the bases of the feathers in adult birds pale or pure white, and in the Andamans there is found the Andaman Jungle Crow (*Corvus coronoides andamanensis*) which is also a larger race with a longer and stouter bill. All the Indian forms, it may be added, are merely races of the Australian Jungle Crow, whilst the form *macro-rhynchus*, under which name the Indian crows have usually been placed, is found in Java. It will be seen, therefore, that, like its grey-necked relation, the Jungle Crow affords an example of species in the making, its representatives in various geographical areas (the Himalaya, the North Indian Plains, South India and Ceylon, Andamans, Australia) having acquired in each area a characteristic facies without having as yet overpassed the rather vague limits which bound the sum-total of characters constituting a species. Although this bird is not a regular migrant and seems to keep to restricted areas in its wilder haunts, yet the races which have taken to scavenging cities and villages for food probably travel over very wide areas in the non-breeding season, and the result of this habit is that the limits of the geographical areas tend to overlap.

THE JUNGLE CROW

3

Except for its preference for a country life, in habits the Jungle Crow is much like the House Crow. Not being such a confirmed



Jungle Crow.

parasite of man, however, it is not so bold as its grey-necked relative. The subspecies *levaillanti*, however, in particular, seems to be in process of semi-domestication and is usually found within easy



distance of human habitations where an easier living is obtainable, either by open scavenging or by following agricultural operations such as ploughing or irrigating. When such are going on, this crow soon discovers the fact and congregates, with many other birds, following the plough and picking up insects exposed by the turning of the soil, and also feeding on insects driven out by irrigation-water. This crow also attends cattle, picking ticks off the animals and also feeding on dung-beetles attracted to their excrement. Other animal food taken under non-scavenging conditions includes caterpillars, beetles, grasshoppers, etc., centipedes, worms, frogs and lizards. The Jungle Crow is very fond of frogs and lizards and, like the House Crow, apparently often worries them to death and then does not eat them. Young chickens, pigeons and even kids are sometimes treated in the same way. The young and eggs of other birds are always eaten if opportunity occurs and this propensity is so well known that, should it venture into or near the tree containing their nests, this crow is vehemently hustled away by such birds as Drongos and Black-headed Orioles. Carcasses of animals are fed on and a corpse floating down a river is usually attended by some of these black ghouls. Vegetable food, however, seems to be taken far in excess of animal and many crops are attacked to a destructive extent. Maize, especially in the case of early crops or under garden cultivation, suffers considerably and in Madras this crow also damages ripening paddy. It is also fond of ground-nuts and, in many districts where this crop is grown, coolies with slings have to be employed as bird-scarers. When in season, the Jungle Crow is also very partial to the flowers of the silk-cotton tree (*Bombax malabaricum*), to all kinds of fig fruit, and to mulberries. Mr. Mason states that the stomachs, which he examined, almost invariably contained vegetable matter much in excess of any other food material. From an agricultural view-point this crow may probably be classed as of neutral or faintly beneficial value, the damage done to crops by pulling up germinating seeds and eating of ripening grain and fruits being compensated, or perhaps slightly




more than compensated, by the good done by its feeding on injurious insects. In sufficiently insanitary localities this crow does a little good by scavenging but is more of a nuisance around houses by its thievish habits. Like most of its tribe, the Jungle Crow is fond of carrying off any bright articles sufficiently small for it to manage and silverware especially is greatly appreciated, the disappearance of small spoons, etc., being often attributable to crows. On one occasion when I was at Coimbatore, a crow was seen to carry off a tea-spoon from the verandah of a bungalow and fly with it onto the roof; a search on the roof revealed not only the missing spoon, but three others which had been misappropriated in the same way. It is hard to say wherein lies the attraction for glittering objects of this kind, which are of no use for purposes of interior economy, but it is certainly a very real one, and little seems to come amiss, even pieces of ice being carried off at times. Sometimes the object taken is simply carried off and dropped, sometimes it is promptly buried, in which case it may be dug up again and re-buried and is then apparently forgotten. A golf-ball, especially a new one, is often an irresistible attraction, being carried off and eventually dropped at a distance; presumably it is mistaken for an egg. Dewar, however, relates how he once scored off a crow by dropping an "approach shot" onto its back as it was swaggering about in the neighbourhood of the "green". He also tells a story of a rook which attempted to hatch out a golf-ball in mistake for an egg.

The Jungle Crow seems to be much less clannish than his grey-necked relation, which is kept in its proper place when the two species enter into competition for food. It is also less gregarious and in the *mofussil* each pair has often its own particular area from which it excludes all other crows. The breeding season is very variable according to locality, usually from January to April, earlier in the South and later in the North. The nest, which is placed in a high tree, is a well-made cup of small, flexible twigs, compactly intermixed with leaves or softer material, and lined

with grass or hair. The usual number of eggs is four or five, rarely more, the egg being about 40 by 29 mm., and spotted, as shown in our Plate, with dull red or brown on a ground-colour of bluish-green. Like the House Crow, this species is also parasitized by the Koel and by the Muscid fly, *Passeromyia heterocheta*.

In Indian folk-lore the crow often acts as a messenger and is supposed to announce approaching visitors. Crows are also often fed as a propitiation to spirits and, when a child is sick, crows and other birds may be bought by its mother and female relatives and released as a propitiation. The crow is generally regarded as a type of knavery and inferiority,—in short, as a wicked rascal. The local vernacular name for this crow in North Bihar is *Kag*, that for the House Crow being *Kowa*.

The Jungle Crow must not be confused with its near relative, the House Crow, "of whom next," as the genealogists have it.



A Kashmiri Songster.



THE INDIAN HOUSE-CROW

(*Corvus splendens*).

“Je n'aime pas les plaisirs innocents.”

Madame de Longueville.

I have never been able to discover any shred of grace about a crow. And what aggravates this state of things is the imposture of its outward appearance. It affects to be respectable and entirely ignores public opinion, dresses like a gentleman, carries itself jauntily, and examines everything with one eye in a way which will certainly bring on an eye-glass in time, if there is a scrap of truth in the development theory.

The Tribes on my Frontier.

OF the Myna, the subject of a later paper, it has been said that it needs little introduction, but the subject of our present paper needs none at all, as it positively thrusts itself into notice and, especially in towns, is the most familiar (in the literal sense of this word) of all Indian wild birds. From early dawn to dewy eve the House-crow is with us, resting somewhere handy to swoop down on any edible morsal and making day hideous with its raucous voice as it relates its misdeeds, which are as black as itself. One feels tempted to stop here and leave our Plate to speak for itself. Is not the House-crow at hand to speak for himself? *Si exemplum requiris, circumspice*. And has he not been treated of at considerable length by many able writers on Indian Birds—Lockwood Kipling, Finn, Cunningham, Aitken, Dewar, Stuart-Baker and others? Dewar has even devoted a whole volume to “The Indian Crow his Book,” which is a signal mark of esteem, as few wild Indian animals have been so honoured—indeed, I can only think at the moment of two others, the tiger, on whose life-story Eardley Wilmot wrote a book, and the sperm-whale, whose doings are chronicled in Melville’s




Moby Dick and Bullen's *Cruise of the Cachalot*. It seems therefore rather a work of supererogation to add to the literature on this bird nor is it possible to compress into a short article the mass of material available.

The House-crow needs little description, being deep glossy-black except for the nape, ear-coverts, the whole head, upper back and breast, which are light ashy-brown. The black portions of the plumage are highly resplendent, when seen at a proper angle, with purple-blue and greenish reflections, whence apparently the specific name *splendens*. The contrast between the paler and darker portions of the plumage varies considerably according to locality and this crow has been divided into subspecies on this character, the Sind House-crow (*Corvus splendens zugmayeri*), found in the extreme North-West of India, having the paler parts almost white in sharp contrast to the black, the Burmese House-crow (*Corvus splendens insolens*) found in Southern Burma, having the dark parts shading into the lighter, which are ill-defined and of a blackish-grey, and the Ceylon House-crow (*Corvus splendens protegatus*), confined to Ceylon, having the light parts shading into the darker but easy to define, whilst the typical race (*Corvus splendens splendens*), shown on our Plate and found throughout the remaining portions of the Indian Empire, has the difference between the grey and black portions of the plumage well-defined. These races, however, tend to run into one another so that individuals from the south of Travancore, for example, are much darker than Northern Indian birds and show a distinct approach to the subspecies found in Ceylon.

The House-crow has little use for a rural life and is scarcely ever seen in the jungle, but any collection of human habitations is sure to have its quota of corvine vagabonds, and, the larger the town, the greater the number of crows. Dewar states that "probably over a million crows dwell in Calcutta" and such a number certainly does not seem exaggerated. About twelve years ago a terrific

hailstorm in Calcutta killed them off in tens of thousands so that the *maidan* was littered with crow corpses, but nowadays the Calcutta crow population seems to be larger than ever. Probably much the same might be said of all large Indian towns. As Aitken puts it, "The crow is a fungus of city life, a corollary to man and sin. It flourishes in the atmosphere of great municipalities, and is not wanting in the odorous precincts of the obscure village innocent of



House-Crow.

all conservancy." Kodaikanal is the only town known to me where crows are not. In large towns, where natural enemies are few and food plentiful, the crow has an easy life, increases in numbers and waxes impudent, but in the *mofussil* he has a rather harder time and wears a more subdued look. At Pusa, for example, there are always a few of these crows hanging around the bungalow but they hardly ever make a nuisance of themselves as they do in towns.



The House-crow makes an easy living by its almost omnivorous habits. In the neighbourhood of houses, it is essentially a garbage-eater and a picker-up of unconsidered trifles in the way of scraps of human food, cooked or uncooked, and preferably acquired by theft. For the crow is not only an outcast scavenger but a thief, delighting to steal from man or dog or other animal or even from its own fellows. Grain, whether in bags in a railway truck or exposed for sale in a shop or in the field, is always an attraction for crows. Fruits, especially figs and mulberries, are eaten greedily and the young and eggs of other birds form especial delicacies, varied by an occasional debauch on carrion. With regard to insect food, crows do not seem to have such a varied diet as one would expect and insects are usually only eaten in any numbers when they occur in swarms; on such occasions, termites, locusts and caterpillars may be eaten to a considerable extent. So that when they eat injurious insects, it is only at a time when the damage caused by these insects has been done and when their destruction even on a large scale makes comparatively little difference to their numbers. Under such conditions an attack on insects by birds is to all intents and purposes useless and of very small value as compared with the good done by an habitually insectivorous bird which feeds on insects throughout the year and which is therefore constantly acting as a check on the increase to undue proportions of its insect prey. A crow only helps to lessen to a small extent the undue proportion of any one species of insect when he notices that they are in swarms and he feels inclined to eat or worry them. For this reason, it is impossible to regard the crow as beneficial from an agricultural view, especially in view of the damage done to grain and fruits. At the best, this bird can only be reckoned as of neutral value. In towns, where the conditions are sufficiently insanitary, it may be considered of some slight value as a scavenger but even against this must be set the annoyance caused by its thievish propensities and noisy habits.



The crow, however, has his good points. His intelligence is distinctly above the bird average and, even when he is making himself a nuisance around houses, a great deal of harmless amusement can often be obtained by watching his wily ways. To vary the poet slightly, very brief observation of his doings soon shows:—

“That for ways that are dark,
And for tricks that are vain,
The House-crow is highly peculiar,
Which the same I am free to maintain.”

A liking for mischief for its own sake seems part of a crow's nature and this may be expressed by pulling up garden plants or digging up newly sown seeds, or even by picking off flowers, or carrying off brightly-coloured objects, or very frequently by taking a delight in worrying other birds and animals for no apparent reason whatever. Robbery is usually at the bottom of this, as when crows worry such animals as kites or dogs, but pure mischief often seems concerned rather than a desire for food. An owl abroad in the daytime is always fair game for crows and a snake which ventures into the open is also often mobbed by crows and other birds such as drongos and mynas. Lizards, such as *Calotes*, are often worried by crows and in his *Tribes on my Frontier* Aitken gives an amusing account of an occurrence of this nature, but a lizard or frog may sometimes be worried to death and then left uneaten, showing that the attack on it is not prompted by hunger. The crow's intelligence is also shown by the occasional holding of *panchayats*, when all the crows of the neighbourhood gather together and vociferously discuss some point of common interest, sometimes apparently the uncorvine conduct of one individual crow who is thereafter set upon by the rest and pecked to death or outcasted. Usually a crow which is damaged in any way is worried to death by its fellows. Yet the crow will certainly rescue a friend in distress if he can and will not hesitate to show his indignation against the cause of suffering to his own comrades. The crow is also a good husband and a tender parent, devoted to his repulsive nestlings,



in whose defence he will not hesitate to attack even man at times.

The House-crow nests at the beginning of the hot weather. In Calcutta building commences about February, but in some parts of Bengal it may even commence in December. The breeding season varies considerably according to locality. Generally speaking, March to May are the favourite months, later in the North and earlier in the South. The nest is a large and untidy mass of sticks, lined with smaller twigs or grass or almost any other softer material, and placed in the fork of a branch of any convenient tree near human habitations. If available, curious materials may sometimes be used for building. Several cases have been recorded in which crows' nests have been built of short lengths of wire and in Bombay one pair of crows constructed a nest of gold and silver spectacle frames of an estimated value of four hundred rupees. Another nest in Madras was made of bits of tin snippings taken from the Tin Bazaar. Both the parent birds collect the material for the nest but its actual construction is apparently the sole duty of the female, the male transferring to his partner, for fixing in the nest, the material which he brings in in his beak. The twigs used may be collected off the ground, or, if not thus available, be broken off growing branches. When building is going on, the male bird accompanies the female everywhere and will not let her go out of his sight. The labour of collecting and building is varied by brief intervals of connubial affection which takes the form of head tickling, but this is not allowed to delay the completion of the nest, which is usually finished in about two days. Thereafter the hen lays in it four or five (sometimes two or three or six, or rarely seven) eggs, which measure about 37 by 27 mm. and vary greatly in colour, the ground being any shade of blue-green, blotched with dull reddish and brown, with small secondary markings, of grey or neutral tint, irregular in shape and scattered profusely over the whole surface of the egg.



At this stage, the crow is often parasitized by the koel (*Eudynamis scolopaceus*), the male of which is a black bird with a very long tail and looking like a drongo, whilst the female koel is glossy black-brown, spotted and barred with white. Both combine to achieve their end of depositing an egg in the crows' nest, and they do this in a very interesting way. Crows seem to hate the sight of the koel, which takes advantage of this fact. The male koel flies up to the crows' nest and attracts the crows to chase him away, a pursuit in which he is quite safe as he is faster on the wing, the koel taking care to lead the crows after him as far as possible. Meanwhile, as soon as the coast is clear, the female koel slips into the deserted nest and deposits an egg, after which she flies off again with a triumphant cry of "*Kuil, Kuil, Kuil*," as a signal that the deed has been done, whereupon the male koel shakes off his pursuers and leaves them to return to their nest. Whether the female koel destroys any of the crows' eggs before depositing her own is not certain but this is probably the case. In any case, the crows do not detect the difference, as they seem to be incapable of counting. Whether the female koel actually lays the egg in the nest is also uncertain, but it is more probable that she lays it elsewhere and then transports it in her mouth until she obtains access to the nest.

Apparently the crows are so pleased with themselves at the successful hatching of the eggs that they are indifferent to the nature of their strange nestlings. At any rate, they appear to be fond of their foster-children. Aitken relates that on one occasion he saw "a pair of crows feeding a clamorous young koel, together with its foster-brother, their own child. It was hungry and clamorous too, but the koel appeared to be the favourite with the parents." The newly-hatched crow nestling is such an abominably ugly little brute that perhaps it is excusable to suppose that even its own parents may prefer a young koel. Be this as it may, they certainly watch over their young with assiduous care and the nest is never left unguarded



for a moment, each parent taking its turn to go out and forage. As Dewar puts it:—"The reason for its devotion is not far to seek. It is the penalty of wickedness. It is a case of thieves knowing the ways of thieves. Crows are notorious robbers of nests; neither eggs nor young birds come amiss to them. They know the evil that is in the corvine heart; hence the careful guarding of the young." But even the watchfulness of the parent birds is unavailing to keep away the fly (*Passeromyia heterochaeta*) which commonly lays its eggs in crows' nests and whose grubs suck the blood of the young nestlings.

The House-crow needs no protection but is quite capable of looking after itself. Hence it receives no benefit in any Province under the Wild Birds' Protection Act, and in most large towns in India it will doubtless be considered that his numbers might be reduced with advantage. Yet other countries have endeavoured at times to acquire him by compulsory immigration. Thus, Finn states that "he has of late years been introduced as a scavenger into Zanzibar," and in December 1902 a consignment of House-crows was sent from Ceylon* to the Malay Peninsula to help in reducing an outbreak of caterpillars on Coffee in the Straits.

The habits of this bird would fill a book and we have only space to refer to its gregarious habit of roosting by night in favourite trees during the non-breeding season. Sometimes, the trees selected as a dormitory serve the same purpose during the daytime for flying foxes. Thus, on the west coast of Ceylon, about thirty-five miles South of Colombo, lies the small island of Barberyn, covered with coco palms, whose tops provide a resting place for these large bats in the daytime and for crows at night, and in the early morning and evening may be seen the passage across the strait dividing the island from the mainland of immense flocks of crows and flying foxes, the one starting out to forage and the other intent on rest.

* See *Spolia Zeylanica*, Vol. I, pp. 23-33, Fig. 13 (1904).



THE BENGAL TREE-PIE

(*Dendrocitta rufa vagabunda*).

A pye sat on a pear-tree,
 Heigho ! heigho ! heigho !
And once so merrily hopped she,
And twice so merrily hopped she,
And thrice so merrily hopped she,
 Heigho ! heigho ! heigho !

Old Nursery Rhyme

THE true Crows, as anyone may observe, have tails which are much shorter than their wings, but many members of the great Crow family have tails much longer than their wings, and this latter group includes the Magpies and Tree-Pies, of which a dozen species occur within our limits, mostly in the Hill Districts of Northern India. The Tree-Pie, however, is rather a bird of the Plains, where it is sufficiently common to be a familiar object in most large gardens, although its curiously metallic cry, rather like the loud squeak from a rusty gate-hinge, if one may imagine a mellifluous squeak, is usually more evident than the appearance of the bird itself. Its most usual call is a sound which may be written *kok-li, kok-li*, but it has a great variety of notes, many of them charmingly melodious in character, others merely hoarse chattering volleys of sound. So far as appearance goes, the Tree-Pie can hardly be mistaken for any other bird found commonly in the Plains, being about eighteen inches long, of which two-thirds is tail, the bill black, the head, neck and breast sooty-brown, the body chestnut-reddish, with some silver-grey on the wings, and



the long tail greyish, darkest at the base and broadly tipped with black. During flight the tail is spread out and, as the tail-feathers are unequal in length, the middle feathers being the longest and the others decreasing in length to the outer pair, which are only about half the length of the middle ones, the expanded tail gives this bird a curious appearance when on the wing. Like many other birds, the Tree-Pie has split up into several local races, of which five have been given subspecific names. Of these, however, three, distinguished by the blending of the colours of the head and back, are confined to Burma and need not be more than alluded to here. The two Indian forms, in which the colours of the head and back contrast strongly, are the Indian Tree-Pie (*Dendrocitta rufa rufa*) which is lighter both above and below and not nearly so richly coloured as the Bengal Tree-Pie (*Dendrocitta rufa vagabunda*) which is darker and more richly coloured both above and below. The former race is found throughout the whole of Southern India as far North as Orissa and in Sind, Afghanistan, and the Punjab as far as Garhwal. The Bengal race occurs from Garhwal to Eastern Assam, throughout the United Provinces, Bihar and Bengal, and is the form figured in our Plate. The habits of both these Indian races, however, are quite similar and both may be considered together under the general name of the Tree-Pie.

The Tree-Pie is a bold and intelligent bird, which goes about in pairs or in small parties, flying from one tree to another and continually prying about for insect or vegetable food. In the early morning especially it is often seen in trees on the hunt for insect food; it is a good climber and supports itself with its claws and tail, rather like a wood-pecker, on vertical or even overhanging stems and branches whilst it searches the crevices of the bark for small insects. Some of its food is obtained on the ground, but most in trees and bushes, and a very small proportion on the wing. The late C. W. Mason stated that "this bird is to a very large extent



Tree-Pie.



a vegetable feeder, though it does not apparently damage crops or planted seeds. It takes a variety of weed seeds and fruits of all kinds including all the common species of *Ficus*, ber fruit (*Zizyphus jujuba*), mulberries, *Sissu* seeds, etc. Of cultivated fruits, when they are in season, it takes peaches, loquats, plantains, etc., and besides eating the fruit on the trees it will often knock off a considerable amount more. Not only does it thus damage the fruit, but it also breaks off small branches (which often contain fruit buds) of brittle-wooded varieties of trees when it alights on them, and is therefore not to be desired in a carefully kept orchard. Leaves and buds of various sorts are also eaten, but apparently only of wild plants. The Tree-Pie's insect food is very varied, but undoubtedly some preference is shown to caterpillars, principally Geometrids and some other smooth varieties—I have never known it touch a hairy one—to beetles, which are mostly Tenebrionids, and to a less extent to the common wasp, *Polistes hebraeus* It does not as a general rule take crickets." It is fond of silk-worm caterpillars and, when it can obtain access to these, may be a nuisance to silk-worm rearers. The food, however, is very varied and one's impression is that it is more largely of an animal nature than Mr. Mason's records indicate. Lizards and spiders are greatly relished and a Tree-Pie will often make a regular practice of hunting around the verandah of a bungalow in the early morning to snap up any lizards or spiders which may be recovering from a surfeit on the insects attracted to the lights the night before. Mr. D'Abreu examined at Nagpur a bird whose stomach contained a mouse, a Buprestid beetle, a caterpillar and two Pentatomid bugs, and at Pusa I have seen one carrying a very fair-sized snake which I managed to make the bird drop and found it to be a *Tropidonotus stolatus*, upwards of two feet long; the snake when rescued was alive and active but bore marks of the bird's mandibles and would undoubtedly have been eaten. The Tree-Pie is also a confirmed robber of the nests of other birds, especially of doves, stealing and devouring the eggs and young of all the smaller birds. Like many



other birds, the Tree-Pie has his good and bad points, but on the whole it is apparently beneficial.

The breeding season is from February to July, from February to March in the South and from May to July in the North, but is not well defined, and eggs may be found both later and earlier than the period normal to any particular locality. When nest-building commences, the parent birds are unusually noisy and the presence of a very voluble pair of birds at this time of year is generally a sure sign that they are beginning to build, and the site of the nest can be located by watching the birds which also resent the presence of other birds on or near the tree which they have selected for themselves. The nest is usually placed well up in a large tree but may at times be built in a thorn hedge, thorny bush or in a Cactus clump. A tree with thick foliage is a favourite site, the nest being well concealed as a rule and never obtrusive to the eye. The nest is a not very large but rather untidy mass of twigs, roots and miscellaneous material, carelessly interwoven and lined with roots or at times with softer material; it is usually about seven inches across, the egg-cavity being about five inches in diameter and usually shallow but at times three or four inches in depth. Three to five eggs are laid as a rule, most often four, rarely six, but in the South only two or three, and the eggs, which average about 29×22 millimetres, belong to two types, one pale greenish blotched and spotted with fuscous, the other pale reddish-white or salmon-colour with blotches of reddish and dark brown and other underlying blotches of lilac and neutral tint; the former type is the commoner in North Bihar. The same peculiarity of producing more than one type of egg is found in other birds but its cause is not very evident: it has been suggested that such variation may be due to the age of the bird, but this hardly appears to be a likely explanation, as in North Bihar, as noted above, the greenish type is the commoner, whilst in the United Provinces the salmon-coloured type preponderates. Both parents assist in nest-building and in feeding the

young and both are also said to take part in incubating the eggs. The young birds are fed almost entirely on caterpillars and perhaps also on fruit to some extent.

The Tree-Pie has not been given legal protection in any part of India. Apparently it is considered well able to look after itself. Being conspicuous, it rejoices in various names in different parts of the country; Stuart-Baker states that the Bengal race is called Bobalink by Europeans, but this name belongs rightly to an American bird and I have never heard it used in India, although it is to some extent descriptive of the Tree-Pie's note; in North Bihar the local vernacular name is *kokaya*, in Bengal it is also called *kotri*, *takka-chor* and *handi-chacha*, in North Cachar *kash-kurshi*, in Assam *khola-khoa*, in Hindi-speaking areas *maha-lat*, at Lucknow *mutri*, in Sind *mahtab* and *chand*, and in Telugu-speaking districts *gokurayi* and *kond-kati-gada*. It will be noted that many of these vernacular names are also expressive of the various calls uttered by this bird.



Indian Magpie.



THE BENGAL JUNGLE-BABBLER OR "SEVEN SISTERS"

(*Turdoides terricolor terricolor*).

THE subject of our present article rejoices in several names. In the new edition of the *Fauna* volume on Birds it is called the Bengal Jungle-Babbler (*Turdoides terricolor terricolor*) whereas it was referred to in the former edition as the Jungle-Babbler (*Crateropus canorus*), and it is also known as the Bengal Babbler, the Seven Sisters, *Chatarhia* (in Bengal), *Pengya-maina* (in the United Provinces), *Kutch-batchia* (in Bihar), *Pedda-Sida* (in Telugu) and as *Sat Bhai*, *Jangli-Khyr* and *Ghonghai* in Hindustani-speaking areas. Of these names, the popular names, "Seven Sisters" or *Sat Bhai*, and the scientific name, *Crateropus canorus*, are the best known applied to this bird, which is sufficiently common everywhere to be a familiar object in all shrubberies of compounds in Northern India north of a line drawn roughly from Orissa to Bombay. As its specific name implies, it is an earthy-coloured bird, a very little larger than a myna with a long tail and a generally untidy look about it, which goes about in small flocks in shrubby places, hopping along over the ground and turning over dead leaves in search of food. In Southern India it is represented by the Southern Jungle-Babbler (*Turdoides terricolor malabaricus*) which differs from the northern bird in being much darker both above and below, most noticeably so on the chin, throat, breast and flanks. The feathers are also darker edged, which shows up more vividly the central pale streaks on the shaft of the feather, these



streaks being indistinct in the northern race. In North-Western India there is a third race, the Sind Jungle-Babbler (*T. terricolor sindianus*), which is paler than the Bengal race, with the upper parts greyer and with the shaft-streaks on the feathers ill-defined or obsolete. The general appearance and habits of all the three races, however, are pretty much the same and the following remarks may be taken to apply to all of them although our Plate represents the Bengal race.

The "Seven Sisters" is especially fond of gardens, being found wherever cover in the shape of a shrubbery is provided. In such places an observer will soon come across a small party of rather bedraggled-looking dusty-brown birds, rustling about in long hops amongst the dead leaves and keeping up a ceaseless gabble of conversation as they follow one another about, turning over the fallen leaves and twigs and peering and prying beneath them for insects, snails and worms. Whilst busy among the leaves they always seem to have an air of dread of finding something terrifying concealed under them and are constantly leaping into the air and starting backward, and yet all the time seem to be in the height of spirits. Although the flock often consists of about half-a-dozen individuals, the number is not necessarily seven and may at times exceed twenty, and why they should be known particularly as the *Seven Sisters* is not especially obvious. Seven is of course a sacred number since Babylonian times, signifying especially the seven planets (the five planets then known together with the sun and moon), whence the number of days in our modern week, and the number as applied to these little parties of birds possibly signifies completeness. Be this as it may, the number is not always seven, as Sir Edwin Arnold observed when referring to this bird in his *Light of Asia*:—

"The nine brown sisters chattered in the thorn."

As regards the second part of the name, as Stuart-Baker well puts it, "their sisterhood or brotherhood they show by the manner

THE BENGAL JUNGLE-BABBLER OR "SEVEN SISTERS" 23

in which each individual resents any interference from outside to any of the party yet retains full liberty to argue, disagree and fight with any one or all of the other six." In cases where one member of the party has been seized by a trained hawk, the others have been known to rush to its assistance and even to rescue it from the hawk. The bond of union resulting from their social habits seems so strong that on the escape from the cage of one individual of a party kept in captivity it has been known to do its best to get in again without further thought of escape. With regard to this trait Finn remarks that "it may be ungenerously suggested that



"Seven Sisters."

such birds are afraid to go about alone, lest the ribald remarks, made in the security of numbers, meet with a just retaliation at the beaks and claws of outraged bird society; and so it may be, but nevertheless there is a well-spring of sincere sociability under the Babbler's frowzy feathering. On the comparatively rare occasions when my captives were still, they employed themselves in affectionately tickling each other's heads as they cuddled together, and I have even seen one diligently employed in endeavouring to clean the wing of a friend, soiled by the bird lime with which its capture had been effected. At the same time it must be admitted that the



addition to their ordinary diet of table scraps of such a delicacy as a cockroach was apt to produce a sad disruption of fraternal harmony. On such occasions one might see one brother prone in the sand while another, holding his head 'in chancery' with one foot, was punching the same with his beak in a manner calculated to awake grave fears for the integrity of the sufferer's skull when the punishment should be over; and once I saw two birds adherent with bill and claw to one and the same cockroach, which a third was devouring, as neither of the joint owners dared let go his hold."

The development of such a degree of clannishness in this bird is doubtless connected with its extremely weak flight. It seems to take to wing comparatively rarely, and when it does, a certain degree of momentum is attained by a violent beating of the wings for a short distance, after which the flight becomes a gliding skim, their pace flagging and their line of flight sinking rapidly. Should a fairly wide open expanse need to be crossed, this can only be done by climbing a tree on one side high enough to allow for the rapid descent that attends their flight, much in the manner of a flying squirrel, which runs up a tree and casts itself off into the air, volplaning downwards on to another tree at a lower level and then running up that and repeating the performance. I have, however, seen this bird rise from the ground and fly upwards into a tree. When in trees, these birds run along the branches in single file, often hopping over one another as they go, and run up and down the tree-trunks, clinging to the bark with great agility even when the surface is nearly vertical. The prehensile power of their feet is not only of use when climbing but also of assistance in holding food.

The "Seven Sisters," as would be expected from its terrestrial habits, is rather a mixed feeder, subsisting largely on insects, small lizards, frogs, worms, etc., which it finds amongst and under dead




leaves and also on wild fruits which always seem to be picked up off the ground.

The late Mr. C. W. Mason examined the stomachs of thirty-six birds at Pusa and found the contents to consist largely of fig and *ber* fruits mixed with a great variety of insects and weed seeds, with an occasional frog, spider or centipede. The food is obtained in jungly or shrubby places, and cultivated areas or crops seem to be rarely visited and this only when there are large trees or jungle close by. The nestlings are fed principally on caterpillars, with a few beetles and an occasional cricket or grasshopper. From an economic point of view, therefore, this bird may be regarded as beneficial to the farmer.

The generally untidy appearance of this bird seems to extend to its nesting habits, the nest being a loose untidy cup, often composed of aerial roots of fig-trees, and placed about twenty feet above the ground in the boughs of small trees or shrubs. At times, the nest may be placed much lower down. The structure of the nest is often so loosely woven that, when the bird is not sitting, the sky can be seen through it from below. The nesting season is any time between March and September but mostly either just before or just after the break of the South-West Monsoon. The usual number of eggs is four, but as many as seven may be laid, and the egg is of a beautiful blue, intensely glossy, and measures about 25 to 20 millimetres. The Babbler is extensively parasitized by that extremely obnoxious bird the Hawk-Cuckoo or Brain-Fever Bird (*Hierococcyx varius*) whose eggs are very similar in general appearance, but usually less glossy and more elliptical in shape. It is also parasitized by the Pied-crested Cuckoo (*Clamator jacobinus*), just as often as by the Brain-Fever Bird, this Cuckoo also laying an egg very similar in colour to that of the "Seven Sisters". In one nest Mr. Inglis found no less than six eggs of the Pied-crested Cuckoo, along with three eggs of the

Babbler, evidently the produce of several individuals of the former species.

The Jungle-Babbler is protected by law throughout the year in Delhi, the United Provinces, Bengal and Assam, and in Madras in the Shevaroy Hills from February to June only.



Common Kingfisher (*Alcedo ispida*).



THE BENGAL RED-VENTED BULBUL

(*Molpastes hæmorrhous bengalensis*).

Of all the throng, the *bulbul* seems to claim
The dearest place; 'tis such a homely bird,
With such endearing ways, fearless and tame,
And everywhere his cheerful note is heard.

Chenar Leaves.

WITH regard to their popular names which are so well known and firmly established that it is difficult to dislodge them from general use, many animals have acquired titles which are perhaps more descriptive than correct. Such, for example, are the "white ant," which is not an ant and not always white, and the "black beetle," which is not a beetle and not necessarily as black as the popular idea paints it. Many similar instances might be quoted and in this connection it is difficult to resist the temptation to refer to the dictionary which defined a lobster as "a little red fish which swims sideways"; whereas of course a lobster is not a fish, it is not red until it has been boiled, and it does not swim sideways. Owing to a similar confusion of ideas or terms the Indian Bulbuls have achieved a somewhat spurious reputation as exquisite birds of song. Both in Eastern and in European poetry the bulbul is frequently referred to as a delightful singing bird, and the dweller in India may well wonder why the Indian Bulbul does not live up to its reputation. The fact seems to be that Indian Bulbul is not the same bird as the bulbul referred to in Persian poetry as the lover of the rose and which is really a nightingale. Our Indian bulbuls have cheerful notes but they are not exactly nightingales.



According to the "Fauna" volume on Birds, some fifty bulbuls are found within Indian limits, of which nine species belong to the genus *Molpastes*. The first six of these, however, are now usually placed together as representing a single species, *Molpastes hæmorrhous*, divided into several geographical races, of which *Molpastes hæmorrhous hæmorrhous* is found in Ceylon and Southern India to about 20° North latitude; *Molpastes hæmorrhous pallidus* takes its place north of this latitude and extends to Bihar and Western Bengal, Rewah, Cutch, etc.; *Molpastes hæmorrhous burmanicus* is found from Manipur to Burma, southwards to Rangoon and eastwards to the Sittoung River; *Molpastes hæmorrhous nigripileus* occurs east of the Sittoung River in South Burma to the Malay Peninsula; *Molpastes hæmorrhous chrysorrhoides* is found in the Kachin Hills, Shan States, and North-East Tenasserim, extending into China; *Molpastes hæmorrhous bengalensis* occurs in the Himalayas, from Kumaon eastwards to Eastern Assam, in North Bihar and Eastern Bengal; and *Molpastes hæmorrhous intermedius* is the race found in the Punjab, North-West Frontier, North Oudh and North-Western Himalayas. In some localities these races run into one another and it seems unnecessary to dilate further on the distinctions between them. The bird shown in our Plate, about half as large again as a sparrow, but with a longer tail, a pointed black crest on its head, and a patch of red feathers beneath its tail, is sufficiently familiar not to require a long description, and may reasonably be set down as a Red-vented Bulbul in one of its numerous forms. It is merely necessary to point out that in other parts of India the local form of this bird may differ slightly from our Plate in the definition or extension of the black on the crown of the head, in the colour of the ear-coverts, chin, or throat, although it always retains the blood-red patch "in the seat of its trousers," as the inimitable EHA so well put it.

These birds are more often found in gardens than in the open country and usually occur in pairs, although sometimes parties of



half-a-dozen or more may be seen together. They frequent trees and bushes and are rarely seen on the ground, for progress on which their short legs are not adapted. In the evening they often take up a perch on a twig at the top of a low tree and thence make short upward flights into the air, returning again to the same perch; Cunningham states that such flights "at first suggest the pursuit of some flying insect, but . . . in reality (are) merely the expression of exuberant nervous energy that is worked off by active exercise and the utterance of pleasant little songs." This may be so at times, but I incline to think that the capture of supper in the shape of some small insect on the wing is usually the object of these short flights, as many small insects are flying at that time and other birds, such as drongos, may be seen catching them at the same time and in the same way, although it is difficult to see what it actually is that they catch. In one of his poems, Sir Edwin Arnold writes of

"The Bulbul, which did chase the jewelled Butterflies,"

and certainly at times the Red-vented Bulbul may be seen to catch and eat butterflies, principally species of *Catopsilia*, and, although its diet is a varied one, insects form a large proportion of it. The late C. W. Mason investigated the stomach-contents of thirty-seven birds at Pusa and found them to contain 129 insects, of which 96 were classed as injurious and 30 as of neutral value. Mr. D'Abreu also at Nagpur found the diet to be a mixture of vegetable matter and insects and observed these birds feeding on long-horned grasshoppers. The vegetable food eaten is largely composed of wild fig fruits at Pusa but is varied according to what is available locally. It is noteworthy that these birds have been credited by several observers with a distinct preference for fruits of a red colour. Cunningham, for instance, says: "Among the fruits that they have a great liking for are those of various gourds, particularly one with beautiful, bright red pulpy fruits" and at Faridpur, in Bengal, this bird has been noted as committing "great havoc in gardens amongst tomatos and chillies, the red colour of which seems to attract them." *Lantana* berries form another class of food of which



they are exceedingly fond and the bird shown in our Plate is depicted sitting on a *Lantana* bush, and in districts in which *Lantana* occurs as a pestilential weed its spread is helped very considerably by the dispersal of the seeds eaten by bulbuls. Bulbuls sometimes do some damage by attacking fruits in gardens and orchards, but the good done by destroying insects throughout the whole year must be offset against this.

The nesting-season varies considerably according to locality, as might be expected in the case of a bird so widely distributed, but in the Plains it breeds chiefly in May, June and July, although a few eggs may be found earlier or later. In Southern India the period is about three months earlier than in the North. The nest, which is placed in almost any sort of tree or bush, is neatly cup-shaped and usually composed of dry grass-stems, very small twigs or stems of small plants, lined internally with fine roots or grass or hairs. Spiders-web is sometimes woven into the outside of the nest. Three eggs are usually laid, but sometimes four are found. The egg varies considerably in size, from about 20 to 27 mm. in length, and from about 15 to 19 mm. in breadth, and is pinkish or reddish white, blotched, streaked or speckled with various shades of red, brownish or purplish red.

The bulbul is comparatively an easy bird to keep in confinement, as its diet is a mixed one. It is often kept as a pet by Indians and it is not unusual to meet a proud owner going for a stroll, carrying a bulbul on a little crutched stick which, in the case of wealthy people, is sometimes made of jade or one of the precious metals. We regret to add that one of the attractions of the bulbul as a pet is its ready pugnacity, which is accentuated, when it is desired to make two birds fight, by starving them beforehand and then showing both a morsel of food, whereupon, as a hungry bulbul will naturally resent competition regarding food-supply, a fight is apt to ensue.



THE BLACK DRONGO OR KING CROW

(*Dicrurus macrocerus*).

THE Black Drongo or King Crow is one of our most common and familiar birds, occurring especially in all cultivated areas and being fond of perching on any suitable upright twig or other support from which it can swoop down to secure its prey, either on the wing or on the ground. The prey consists almost wholly of insects, and practically wholly of injurious insects, so that this bird is most distinctly a valuable ally of the farmer and deserves every encouragement and protection.

The Drongos form a family of birds well defined by their black plumage and long forked tail composed of ten feathers. Less than twenty species occur within Indian limits and these are divided into about seven genera, again subdivided into two groups according to the relative lengths of the outer tail-feathers. In one group, in which these feathers exceed the middle ones by at least twice the length of the wing, we may briefly notice the larger Racket-tailed Drongo (*Dissemurus paradiseus*), which occurs in practically all the more hilly parts of India and Burma, is black, glossed with blue, with a tuft of feathers on its forehead, and which has a really fine song and, according to Oates, is perhaps the best singing-bird of the East. In the other group, characterized by the outer tail-feathers not exceeding the middle ones by so much as the length of the wing, we have the genus *Dicrurus*, with an untufted forehead and a deep bill, comprising about eight Indian species, amongst which *D. macrocerus* is distinguished by its entire plumage being a deep glossy black and its outermost



tail-feather exceeding the middle pair by a distance greater than twice the length of the tarsus. The Black Drongo is the commonest bird seen near dwellings in the Plains and any jet-black bird, about the size of a bulbul, with a long forked tail, seen in cultivated areas in the Plains, is likely to be this bird. The whole plumage is deep black, everywhere glossed with steel-blue, the latter character being perhaps rather accentuated in our Plate, which shows a small white spot which is sometimes present at the angle of the gape. Young birds have the wings and lower plumage brownish mixed with white, and very old birds may be entirely black.

The specific name *ater*, under which the Black Drongo has hitherto been referred to as a rule in Indian literature, of course means black, in reference to the colour of this bird, but it is a feeble name to have been bestowed on it, and a name such as *audax* would seem better suited to a bird of this character. Its alternative name of King Crow is probably derived from the fierce manner in which it attacks and drives away any crows which approach near its nest during the breeding season. Except for its colour it is not itself crow-like either in build or disposition. In its nature, indeed, it partakes of a good deal of the piratical, watching from its perch above until some industrious Hoopoe or Myna has found a tasty morsel of insect food and then swooping down and robbing the rightful owner with a dexterity evidently only acquired by a regular course of original sin in the line of highway robbery. The victim, if it be a Myna or similar bird, sometimes shows fight or endeavours to escape with its prize, but the Hoopoe is more of a philosopher and allows itself to be robbed almost with impunity, merely ruffling up its feathers and then starting in again to peck away in the ground in the search for another grub. In any case the result is usually the same, and the Drongo secures the spoil and flies off with it to its perch. This little comedy may be seen enacted daily in almost any Indian garden, especially when the Hoopoe has

young and is collecting food to take back to the nest, and it is noticeable that the Drongo only levies just enough toll not to discourage its victims unduly and drive them away altogether.



Drongo.

The Drongo, however, is by no means wholly a parasite, living at other birds' expense, but catches for itself the larger proportion of its food. The late C. W. Mason examined the contents of twenty-seven adult and four young birds at Pusa and Mr. D'Abreu has recorded the contents of seven birds at Nagpur, and these records

show that the food consists entirely of animal matter, practically wholly of insects and in an overwhelming proportion of injurious insects, such as crickets, grasshoppers, moths, bugs, and insect larvæ. The prey is captured either on the wing or on the ground, in which case the bird swoops down upon it and either eats it forthwith or flies off with it to its perch. Mr. Mason gives a long and interesting account of the feeding-habits (*Ind. Agric. Entl. Mem.*, Vol. III, pp. 69—79 ; Jan. 1912), which we need not reproduce here. The Drongo is one of the birds which accompany cattle, often perching on their backs, but apparently rather for the sake of the grasshoppers and other insects disturbed by the movements of the animals than for the sake of the flies which annoy them. It is also a regular visitor to ploughing operations, perching on upstanding clods or other convenient vantage-spots, and picking up cutworms and other insects exposed by the plough. As Mason notes, however, "he is by no means active on the ground under the best circumstances," his long tail, which serves so well as an organ of flight, seeming to get in the way when on the ground. It is noticeable that the Drongo never hunts on the ground for insects, but always watches from some convenient perch, swooping down to capture insects seen on the ground. The Drongo is naturally a pugnacious and fearless bird, attacking without hesitation birds much larger than itself. At the breeding-season its natural pugnacity is much augmented and it drives away any birds, especially crows, which dare to approach the tree in which its nest is placed. As noted in another place, the Indian Oriole frequently builds in the same tree as the Drongo, the latter thus serving to protect the Oriole's nest as well as its own, and at the moment of writing these lines there is within a few yards of my bungalow a *sissu* tree in which a pair of Drongos and of Indian Orioles have each a nest quite close to one another. A pair of Jungle Crows commenced the nesting-season by building in the topmost branches of the same tree, but now they dare not approach it and probably the young crows have flown by now in



any case. Drongos sometimes fight amongst themselves and I have seen a couple so intent on squabbling on the ground that a jackal was able to run up, knock them both down with its forepaws and carry them off.

The nesting-season of the Black Drongo is from April to August but May is the usual time in Bihar. The nest, which is a broad shallow cup, is usually placed fairly high up in a tall tree in some fork of a branch not quite on the outside of the tree ; it is composed of small twigs and stems and roots of grass, neatly and tightly fastened together, and bound around on the outside with more or less cobweb, in which a few feathers are sometimes entangled. The cavity is broad and shallow, about four inches broad by one and a half deep, and lined with horse-hair or fine grass stems or roots ; the bottom of the nest is very thin, but the sides are rather thick and firm. Four eggs is the normal number, but there may be three or five at times. The egg is either dull pure white without markings or pure white sparsely spotted with deep blackish brown, or pure creamy pink blotched with reddish-brown to deep salmon-pink blotched with claret colour ; it is noticeable that all the eggs of one clutch nearly always seem to belong to one or other of these types but Mr. C. M. Inglis has found a clutch with two white eggs and one spotted one ; the average size of the egg is about 25 mm. long by 19 mm. broad.

The Drongo is protected by law throughout the whole year in Delhi, the United Provinces, Bengal, Assam, and Burma. It is, as already remarked, a most useful bird which deserves every protection and encouragement. The latter aim can be attained, in the case of cultivated areas, by the provision of suitable perches for the birds to rest on.

This Drongo has long been known to Indian ornithologists as *Dicrurus ater* but its correct name seems to be *Dicrurus*



macrocerus, Vieill. ; it is divided into four races, of which the typical one, *D. macrocerus macrocerus*, ranges throughout India to the foothills of the Himalayas ; the Himalayan form, *D. macrocerus albirictus*, ranges from the Himalayas to the Northern Chin Hills ; the Burmese subspecies, *D. macrocerus cathoecus*, is found in Burma, Siam, and South China ; and the southern form, *D. macrocerus minor*, occurs in Ceylon and doubtfully in Travancore.

It is on the wing until well into the dusk and is one of the earliest birds to call in the morning and may at times be heard throughout the night. It has a variety of notes and is an excellent mimic. Donald records that he has heard one imitate a *shikra* (*A. badius*) so as to frighten away a pair of Mynas and thus secure a worm which they had caught.

Mr. C. M. Inglis has seen one chase a Laggar Falcon (*Falco jugger*), but in this case the tables were turned, for the falcon stopped in its flight and the Drongo shot ahead and was captured by the falcon. The smaller bird in our plate is shown attacking one of these falcons which had approached its nest, shown on a tree in the background.

Although the Black Drongo is so pugnacious and intelligent, it is sometimes imposed upon by the Drongo Cuckoo (*Surniculus lugubris*), a bird which is itself most remarkably like a Drongo in general appearance and which lays its eggs in Drongos' nests.



THE TAILOR-BIRD
(*Orthotomus sutorius sutorius*).



THE TAILOR-BIRD

(*Orthotomus sutorius sutorius*).

No better a musician than the wren.

Shakespeare—*Merchant of Venice*.

Natura maxime in minimis miranda.

THE Warblers include a very large number of kinds of small birds which, with few exceptions, are plainly coloured and alike in both sexes. Over one hundred species occur within Indian limits, many being migrants, passing the summer in Central or Northern Asia, whilst others are permanent residents in the Plains and lower ranges of the Hills. Amongst these latter are included the Tailor-birds, so called from the remarkable skill which they display in the construction of their nests which are placed in a receptacle made by sewing together the edges of one or more leaves, and of these little birds the most common and widely-distributed is the Indian Tailor-bird (*Orthotomus sutorius sutorius*). This is a tiny greenish-brown bird, whitish below, in general appearance like a wren, but with a decidedly longer tail; if further description is needed, it may be noted that the forehead is reddish, shading off into ashy-grey on the back of the head, and that on each side of the neck there is a short black bar or spot, usually concealed but plainly visible when the bird stretches its neck to utter its note, which it is doing continually. Indeed, as Cunningham has so aptly put it, "whenever memory reverts to the experiences of summer in the Plains of India, it can hardly fail to recall the loud shouts of the Tailor-birds, as they travel about ceaselessly among the shrubs. Even at those times of day when the breathless heat



and cruel glare have reduced almost all other birds to relative silence; when even the crows sit about in pairs in the shade, gasping with widely gaping bills and incapable of anything beyond whispered conversation, and when the still and fiery air is only rarely disturbed by the querulous whistle of a kite, even then the Tailor-birds are all alive with noisy excitement. Whilst listening to them, or to the cries of other loud-voiced small birds, one realizes the beauty of the dispensation that has decreed that in the animal kingdom there should be no necessarily direct ratio between size and vocal power; an elephant with a voice on the scale of that of a tailor-bird would have been a nuisance to a whole district! Even the longest use and wont leave it a ceaseless marvel how such pygmies can manage to make such a hubbub, whilst they run and creep about among the bushes, more like little brown mice than any feathered creatures. They have two common calls, the first consisting of an urgent repetition of the syllable 'pēēt,' and the second, even more insistent and sounding, 'pe peep, pe peep, pe peep, pe peep.' Long after most other birds are silent; after even the crows and mynahs have finally settled down for the night, and only an occasional belated kite is audible, their call may still be heard issuing from the flower-beds and shrubberies, where the birds continue to run mouse-like about in the gathering gloom, jumping after the insects lurking among the leaves. When highly excited over anything they shout their loudest, and, with their tails so excessively elevated that they come to point obliquely forwards over their backs, look more like demented wrens than anything else. Whilst engaged in hunting over a shrub they run quickly along the twigs, shouting noisily all the while, and every now and then snatching at insects, and, even when flying, they continue to call aloud with a reckless expenditure of breath." The above has been quoted at length, as it seems to be a charming character-sketch of the little bird in question, which, although inconspicuous and apt to be overlooked, will now perhaps be recognizable to our readers. The typical race, as shown on our



Plate, of the Indian Tailor-bird (*O. sutorius sutorius*) occurs throughout the Plains and lower Hills of India, Ceylon, and North and Central Burma, being replaced in Southern Burma, Siam and the Malay Peninsula by the Malay Tailor-bird (*O. sutorius maculicollis*), which differs by having white streaks on the ear-coverts.

As noted above, Tailor-birds are mainly insectivorous in their diet. The late Mr. C. W. Mason examined the stomach-contents of four birds at Pusa and found them to consist entirely of insect remains, mainly of small beetles, bugs, ants and flies; Mr. D'Abreu found much the same in the case of three birds examined at Nagpur. They seem, therefore, to be useful little birds to have in a garden.

The breeding-season of the Indian Tailor-bird is mainly in June and July, but nests may be found from April until August. The structure and design of birds' nests in general may vary from just nothing at all, the eggs being laid on the bare ground, to elaborate mounds such as are constructed by the Bower Birds, but for ingenuity, as applied to its construction and concealment, the Tailor-bird's nest is hard to beat. Pennant seems to have been the first to give any description of it and we reproduce the rather quaint figure given in his *Indian Zoology* (second edition; 1790) where he says that in India "the brute creation are more at enmity with one another than in other climates; and the birds are obliged to exert unusual artifice in placing their little broods out of the reach of an invader. Each aims at the same end, though by different means. Some form their pensile nest in shape of a purse, deep, and open at top; others with a hole in the side; and others, still more cautious, with an entrance at the very bottom, forming their lodge near the summit. But the little species we describe, seems to have greater diffidence than any of the others: it will not trust its nest even to the extremity of a slender twig, but makes one more advance to safety by fixing it to the leaf itself. It picks



Nest of the Tailor-bird amongst mango leaves (Pennant's *Indian Zoology* (1790), Pl. 10).



up a dead leaf, and, surprising to relate, sews it to the side of a living one, its slender bill being its needle, and its thread some fine fibres; the lining, feathers, gossamer, and down. Its eggs are white. The colour of the bird light-yellow: its length three inches, its weight only three-sixteenths of an ounce, so that the materials of its nest, and its own size, are not likely to draw down a habitation that depends on so slight a tenure." So far as it goes, Pennant's account is fairly accurate except in the statement that the bird picks up a dead leaf and sews it on to the side of a living one. The dry leaves which are often found attached to the outside of the nest are accounted for by the fact that these leaves, which have originally been pierced by the bird whilst they are still living and attached to the tree or plant, often die and decay *in situ*, either as a direct result of the injury caused to them or by interference with the free access of air to the tissues.

The nest may be placed at any elevation, such as high up in a mango-tree or low down in a brinjal-plant but more usually they are built comparatively low down, often within a couple of feet of the ground. The kind of leaf selected seems to be rather immaterial, provided that it is fairly large and sufficiently strong to hold the strain of the stitches, and in gardens the leaves of mango, guava and brinjal are often used. As a rule only one or two leaves are used but occasionally, when the nest is placed on a plant with small leaves, such as oleander, a dozen leaves may be fastened together. Generally, the nests hang down more or less vertically, as shown in Pennant's figure, but sometimes they lie almost horizontally with the opening between the lower edges of the leaves, so that the nest is well protected in rainy weather.

The actual construction of the nest has been observed by Mr. A. G. Pinto, as reported by Dewar in his *Birds of the Plains*. In this case the nest was placed in the leaf of a *Dracena* plant growing on a verandah and we cannot do better than quote our



authority for what took place :—" One of the leaves of the plant was so curved that its terminal half was parallel with the ground. Upon this she commenced operations. The first thing she did was to make with her sharp little beak a number of punctures along each edge of the leaf. In this particular case the punctures took the form of longitudinal slits, owing to the fact that the veins of the *Dracaena* leaf run longitudinally Having thus prepared the leaf, she disappeared for a little and returned with a strand of cobweb. One end of this she wound round the narrow part of the leaf that separated one of the punctures from the edge ; having done this, she carried the loose end of the strand across the under surface of the leaf to a puncture on the opposite side, where she attached it to the leaf and thus drew the edges a little way together. She then proceeded to connect most of the other punctures with those opposite to them, so that the leaf took the form of a tunnel converging to a point. The under surface of the leaf formed the roof and sides of the tunnel or arch. There was no floor to this, since the edges of the leaf did not meet below, the gap between them being bridged by strands of cobweb. This was a full day's work She next went on to line with cotton this *cul-de-sac* which she had made in the leaf. She, of course, commenced by filling the tip, and the weight of the lining soon caused the hitherto horizontal leaf to hang downwards, so that it eventually became almost vertical, with the tip pointing towards the ground. When lining the nest the bird made a number of punctures in the leaf, through which she poked the lining with her beak, the object of this being to keep the lining *in situ* All this time the edges of the leaf that formed the nest had been held together by the thinnest strands of cobweb, and it is a mystery how these can have stood the strain. However, before the lining was completed, the bird proceeded to strengthen them by connecting the punctures on opposite edges of the leaf with threads of cotton. Her *modus operandi* was to push one end of a thread through a puncture on one edge and the other end through a puncture on the opposite edge



of the leaf. The cotton used is soft and frays easily, so that that part of it which is forced through a tiny aperture issues as a fluffy knob, which looks like a knot and is usually taken for such. As a matter of fact, the bird makes no knots; she merely forces a portion of the cotton strand through a puncture, and the silicon which enters into the composition of the leaf catches the soft, minute strands of cotton and prevents them from slipping Sometimes the connecting threads of cotton are sufficiently long to admit of their being passed to and fro, in which case the bird utilizes the full length."


It is only the hen-bird which constructs the nest. Although the two sexes are coloured much alike, in the breeding-season the male bird acquires very elongated middle tail-feathers, projecting about two inches beyond the normal length of the tail, so that the sexes are readily distinguishable. We cannot say whether his extra length of tail is an impediment to nest-building or whether it makes him too proud to work or whether he is merely lazy; but to the hen-bird must be given all the credit of the wonderful method of nest building. Aitken, however, in his *Common Birds of Bombay*, implies that it is the cock-bird which builds.

The leaves containing and concealing the nest are fastened together with any suitable material that is available such as cobweb, caterpillar silk, thread, wool, or vegetable fibres. When a nest is being built near a house the bird will often make use of threads of cotton or loose strands from a coir mat. Jerdon says, "I have seen a Tailor-bird at Saugor watch till the native tailor had left the verandah where he had been working, fly in, seize some pieces of thread that were lying about, and go off in triumph with them; this was repeated in my presence several days running."

The nest itself is a neat cup, about three inches deep and two inches in diameter, constructed of wool, cotton, with a few

hairs and fine vegetable fibres, the cavity being always very softly lined. Three or four (more frequently three) eggs are laid, measuring about 16 by 12 mm., the ground-colour being either reddish-white or pale-bluish-green, the former being the more common, but all the eggs in any one nest always belong to one or the other type; the markings on the eggs consist of blotchings and clouds tending to form a bold irregular cap around the larger end, and also of smaller, brownish-red specks and splashes scattered more or less over the whole surface of the shell. When sitting on the eggs the hen-bird lies very close and does not usually fly out until the nest is actually touched or shaken.

The young birds are well looked after by their parents for some time after they have left the nest and small family-parties, consisting of the parents and their young, may often be seen at that time of the year.



A beautifully constructed nest, found in a hedge at Pusa. The outside is covered with small fragments of bark. [N.B. This is, of course, not the nest of a Tailor-bird.]



THE BLACK-HEADED ORIOLE
(*Oriolus luteolus luteolus*).



THE BLACK-HEADED ORIOLE

(*Oriolus luteolus luteolus*).

Yon joyous bird the golden oriole—
Fairy embodiment of living gold,
In melodies so blithe pours out his soul,
And lights the Dal's dim green with colour bold.

Chenar Leaves.

ORIOLES, commonly called "Mango birds" in India, are well-known to all observers of bird life on account of the conspicuous yellow which is the prevailing hue in nine out of our ten different forms, the tenth (*O. traillii*) being black and maroon-red. Of the four species, represented by nine subspecies to which the term Yellow Orioles may be applied, only two are common in the Plains of India and these are the Indian Oriole (*Oriolus oriolus kundoo*) and the Indian Black-headed Oriole (*O. luteolus luteolus*), in both of which species the tail is black and yellow or green but in the former the nape and crown of the head are yellow or greenish-yellow with a small patch of black through the eye, whilst in the latter species these are black, as is the upper breast. Some females are nearly as brightly coloured as the males. Both species are very similar in their habits, being thoroughly arboreal birds, rarely seen on the ground, and feeding mostly on fruit and such insects as are found on leaves and stems.

Cunningham, in his book *Some Indian Friends and Acquaintances*, gives a charmingly written account of the Black-headed Oriole in Calcutta, in the course of which he says:—"It would be hard to imagine any plumage more beautiful than that of the mature



birds with its brilliant contrasts of vivid yellow and shining black and though that of the females and young birds is not so striking, owing to the greenish tone and streakiness of the yellow parts, it has very decided beauties of its own in its delicate gradations and pencillings of colour. They have a truly astonishing variety of notes, almost all of them charmingly melodious in character. As a rule, they go about in pairs, who pass from tree to tree 'crying and calling' to one another at brief intervals. When they are in their very fullest voice the one bird cries, 'Yū, hū a yu,' and the other almost immediately replies, 'Tu hū ēē'; when very much out of voice they often can do no more than cry 'Tehēē' like Alisoun in the *Miller's Tale*; and between these extremes there is a whole range of very distinct calls that only agree in conveying a sense of joy and fulness of life and melodious contentment with it. All of these are highly characteristic and distinct from the notes of any other kinds of birds, save one or two of the most fluty cries of the common tree-pies. It is delightful to see any living things so full of the pure joy of existence as a pair of Orioles always seem to be when they come leaping through the air into a garden, calling as they go; or, after they have alighted in a tree, chasing one another about from bough to bough with their golden plumage shining out among the surrounding green. Now and then a solitary bird will take to haunting a garden for a time, making its appearance regularly day after day at a particular time, in order to visit certain trees and talk softly to itself as it goes on its way; but it is only when in pairs, or in a small family party of three or four birds, such as may sometimes be seen soon after the nesting season, that they fill the air 'with their sweet jargoning.' The solitary birds occasionally seem to be soured by the want of companionship, and travel round hustling other birds and knocking them off their perches out of gratuitous ill-temper—conduct of which paired birds are never guilty. In addition to the manifold modifications of their regular melodious calls, they sometimes utter harshly cawing notes, and the young birds for a time indulge in churring cries somewhat like those of starlings."



During the Rains, this bird often makes a whistling call which may be written as *Whittu*; by mimicking this call it may often be induced to reply for quite a long time.

As noted above, the Black-headed Oriole feeds chiefly on fruit and small insects. The late C. W. Mason examined the stomachs of twenty-three birds at Pusa and found that seventeen of these had fed on wild fig fruits and five of these contained nothing else; the eighteen which had fed on insects contained 95 insects, of which four were classed as beneficial, 73 as injurious and 18 as neutral. Five birds had eaten insects only. In the Central Provinces Mr. E. A. D'Abreu found a Pyralid caterpillar and *Ficus* fruit in the stomach of one bird examined on 24th January 1914. At Pusa also we have watched this bird feeding on masses of a mealy-bug clustered on the stem of a wild vine growing on a *sissu* tree. So far as agriculture is concerned, therefore, this bird may be considered beneficial. It has not been noted to attack cultivated fruits.

Many authors (e.g., Oates and Dewar) note that this bird is strictly arboreal in habits, never descending to the ground; but it is occasionally seen on the ground capturing insects.

The nest is a cup-like structure which is usually placed in the fork of a branch and lashed in position by means of fibres which are wound first around a branch, then passed under the nest, and then wound around another bough. In the case of the Indian Oriole (*O. oriolus kundoo*) Dewar (*Birds of the Plains*, pp. 138-139) notes:—
“A very curious thing that I have noticed about the Indian Oriole's nest is that it is always situated either in the same tree as a King-crow's nest or in an adjacent tree. I have seen some thirteen or fourteen Orioles' nests since I first noticed this phenomenon, and have in every case found a King-crow's nest within ten yards. The drongo builds earlier, for it is usually feeding its young while the Oriole is incubating. It would therefore appear that it is the Oriole which elects to build near the King-crow. I imagine that it does so



for the sake of protection ; it must be a great thing for a timid bird to have a vigorous policeman all to itself, a policeman who will not allow a big creature to approach under any pretext whatever." The Black-headed Oriole, however, does not seem to have been noticed to have police protection in this manner.

The breeding season of the Black-headed Oriole is from March to August. Both birds appear to take part in the construction of the nest which is almost always firmly attached to the extreme terminal twigs of an upper horizontal branch of a large tree, some twelve to thirty feet above ground-level. The nest itself is a deep cup, the egg-cavity measuring about three inches in diameter and two inches in depth. It is composed chiefly of tow-like vegetable fibres, thin slips of bark and the like, and is lined inside with very fine tamarisk twigs or thin grass, and on the outside it is generally more or less covered over with odds and ends, bits of lichen, thin flakes of bark, etc. Three eggs are laid as a rule, but occasionally there may be two or four. The egg averages about 28.5 mm. long by 18 mm. broad, the shell fine and moderately glossy, usually creamy or pinky white with sparse spots and streaks of dark brown and pale inky purple.

The Black-headed Oriole has long been known in Indian ornithological literature under the specific name *melanocephalus*, which is Greek for "black-headed," a name applied by Linnaeus in 1766, presumably to the male bird. The female, however, was described by Linnaeus in 1758 under the name *Sturnus luteolus* and, by the Law of Priority in nomenclature, the name *luteolus* takes precedence as the correct name for this bird. The form occurring in Tenasserim is separated under the subspecific name of *Oriolus luteolus thaiacous*.

This Oriole is protected throughout the year in Bombay, the United Provinces, Bihar and Orissa, Bengal, Assam, Burma, Madras, and Mysore.



THE COMMON MYNA
(*Acridotheres tristis tristis*).



THE COMMON MYNA

(*Acridotheres tristis tristis*).

Striped squirrels raced, the mynas perked and picked.

The Light of Asia.

Talkative Mynas in pairs are pacing the ground, for among the sprouting grass they will find the lubberly sons of the Grasshopper. *Acridotheres*, the grasshopper-hunter, is the happiest of ornithological names; but I never could understand why it should be called *tristis*. Who ever saw a sad Myna?

A Naturalist on the Prowl.

THE Common Myna needs little introduction as it is of common occurrence in every garden throughout India. "Neat but not gaudy" seems to be its motto as it goes through life in its decorous, but by no means dingy, colouring of black, brown and white, finished off with a bright yellow beak and legs and a yellow patch behind the eye. The wings are black, with a white bar which is most conspicuous during flight, when, according to Lockwood Kipling, "a curious effect of rotation" is produced. The head, neck and upper part of the breast are also black, as are the tail-feathers, which have broad white tips also especially conspicuous during flight. The rest of the plumage is brown. Occasionally colour variations occur and albinos are not very rare and Finn records examples with pale cinnamon body-plumage; for some time there was an individual in my compound with a bright yellow head—possibly due to moulting. The two sexes are coloured alike but the hen is slightly smaller, the cock bird having a larger head and more massively built body. The Common Myna, as represented on our Plate and found throughout India and Burma, is the typical form, the Ceylon race (*Acridotheres tristis*



melanosternus), representing a subspecies which differs in being darker and in having the outer webs of the earlier primary-coverts black. South Indian examples are darker than those from Northern India.

As is expressed by its generic name, *Acridotheres*, which means a grasshopper-catcher, the Common Myna is mostly a ground-living bird, pacing about in open spaces, and especially over grassy lawns, on the hunt for grasshoppers and other insects. Like the Cattle Egret, it is a constant attendant upon cattle, following the animals and catching the insects disturbed by their movements. For the same reasons, it is greatly interested in cultural operations, such as ploughing or the cutting or irrigation of crops, as such occasions provide a rich harvest of insect prey. The insects eaten by this bird are practically all of injurious kinds—grasshoppers, crickets, caterpillars and beetle grubs for the most part—but the Myna is a very general feeder and spiders and worms are eagerly sought for and devoured, whilst grain and fruits lend variety to its diet. The fruits eaten are mostly wild figs and no great toll is levied on cultivated fruits. Cereals, however, are sometimes attacked to some extent and some little damage is done at times, especially to maize which is preferred whilst the seeds are still soft and unripe. As in the case of many other useful birds, however, it is necessary to take a broad view and not grudge a small toll levied on cultivation in return for the great assistance rendered throughout the year in reducing insect crop-pests.

In his *Birds of the Plains*, Dewar calls the Myna “a bird of character,” and such undoubtedly it is, standing no nonsense from other birds, whether of its own or of other species. A pair of Mynas, accustomed to quarter a restricted patch of lawn for insect prey, will strongly resent intrusion on their hunting ground by any other bird and will rigorously put into practice an unwritten notice that “Trespassers will be prosecuted.” As Finn puts it, “Bold,



vigorous and pushing, he secures to himself a large share of all good things in the way of insects and fruit that may be going, and is a bird of remarkable all-round abilities, though not particularly graceful in his movements." Few birds dare stand up to a Myna and there are very few birds that a Myna will hesitate to assault. For this reason, the Myna has not proved an altogether unmixed blessing in those countries to which it has been introduced from India as it is too powerful and too free a breeder to be allowed to increase without checks and thus often renders itself a nuisance by its aggressive habits towards other, not less useful but less pugnacious, birds. It has been introduced from India into Mauritius, Australia, New Zealand, and Hawaii and in the last mentioned locality has shown itself a decided nuisance by turning out pigeons. In the Andamans also, whither it was introduced in 1873, it seems to be doing its best to oust the pretty little native white Myna (*Sturnia andamanensis*). It was introduced into Mauritius to destroy grasshoppers and was abundant and perfectly naturalized around Port Louis when I was there in 1905.

In spite of his assertive nature, the Myna is an extremely interesting acquaintance, as his very boldness makes his actions easy of observation whilst his universal occurrence provides endless opportunities of watching his happy habits. Unlike the crow, with its furtive and self-conscious manner, the Myna makes itself thoroughly at home and at its ease around or even in a human habitation, where its actions seem to be always perky and devoid of any sense of shame. Indeed, as Cunningham aptly says, Mynas "always make themselves entirely at home in a house, taking it completely for granted that they are quite at liberty to drop in and stay whenever and for as long as they like." Sometimes, indeed, they make themselves rather too much at home, as when they insist on nesting in some convenient hole in a house and litter the surroundings with a miscellaneous collection of nesting materials. Here again we may quote from Cunningham's excellent account of a



pair of Mynas which helped to while away the hours of his first hot weather in Calcutta :—"He and his wife had elected to place their nest on the cornice beneath the beams in one corner of the open roof of my room, and were constantly coming in with fresh stores of building materials. It was quite refreshing to see the supreme satisfaction that they derived from the progress of their work—a satisfaction that every now and then became so acute as to call for a short rest and jubilant little song. Merely to watch the construction of a myna's nest is a liberal education ; it is like watching the steps in the formation of a local museum. Their taste in materials is so catholic that one never knows what curio may not be brought in. Sticks, straws, feathers, rags, small bones, and pieces of paper are all deemed valuable, and a very special worth would seem to attach to the cast skins of snakes, for, in any case where these are attainable, they are almost sure to be worked into the growing heap of rubbish. The pity is that in their effort to bring in exceptionally bulky materials they are apt to drop them about, and, although snake-skins and feathers may be interesting and even decorative additions to the furniture of a room, great pieces of paper or rag, of unknown origin and very doubtful purity, can hardly be regarded as very desirable additions to one's surroundings."

The nest, as may be gathered from the preceding, is an untidy mass of dry grass and miscellaneous rubbish, usually warmly lined with feathers, and placed in a hole, where this is available, either in a tree or in the wall or roof of a house. Sometimes it may be placed in creepers growing on a wall, or on any sheltered ledge, occasionally in palms or other trees. The nesting season lasts from the beginning of the hot weather until well on in the South-West Monsoon. As a rule, four or five eggs are laid and these are of a glossy bright blue colour, about 30 mm. by 21.5 mm. Mr. C. M. Inglis has on one occasion taken a Koel's egg from a nest of this species which also held three of the Myna's own eggs ; this is, of course, exceedingly unusual. The young are fed




mostly on an insect diet consisting of caterpillars, beetle grubs, grasshoppers and crickets, varied by occasional worms and soft fruits.

Mynas are of course often kept as cage-birds and birds which are reared up from the nest become very tame and fearless. The Common Myna is a good mimic also, although not able to acquire such a wide range of vocabulary as the Grackles or Hill Mynas. Normally, it is rather a noisy bird, its song, to be heard throughout the day, comprising a strange mixture of harsh gurglings and liquid notes. E. H. Aitken described its notes as "Keeky, keeky, keeky.churr, churr, kok, kok, kok" and these sounds are fairly descriptive. As it utters the *kok, kok, kok*, it bobs its head in a characteristic manner. When a small flock is feeding on the ground and one bird sees cause for alarm, it takes to wing with a shrill cry of alarm to its fellows. As a rule, such small flocks seem to consist of two or more individuals, generally a cock and hen and their family. Mr. C. M. Inglis has on one occasion seen this Myna feeding the young of the Pied one. During the off-season for nesting, however, they often congregate in large flocks in the evening to roost in company on favourite trees providing dense leafy cover and at such times the deafening din that arises is quite ear-splitting, as every bird seems to have to argue about its proper perch and then to proceed to gossip at the top of its voice about the events of the past day. During February, when the silk cotton (*Bombax malabaricum*) trees are in flower, like flying-foxes and many other birds, they are very fond of the liquor that is to be found in the lower part of the large red flowers, and the competition for this often creates a prodigious fuss almost equal to that at roosting-time.

The Common Myna has many vernacular names in India. It is known as *Maina* or *Desi-maina* in Hindi, as *Salik* or *Bhat-salik* in Bengali, as *Bemni* and *Salu* in Chota-Nagpur, as *Salonka* in

Mahratti, as *Gorwantera* in Kanarese, as *Goranka* and *Gorinka* in Telugu, and *Zayet* in Burmese, while in the Central Provinces there is a local name *gulgul*, of which *gural* in the Punjab is apparently another form. In Hindu mythology the Myna is sacred to the God Ram Deo and sits on his hand. In local stories and talk this bird is also a favourite.

The Common Myna is protected by law throughout the year in Bombay, the United Provinces, Delhi and Bengal, and in Burma for trade purposes in reserved forest areas. In Mysore it is presumably protected under the heading, "birds of song, absolutely." On the whole, in India at all events, it seems to be a decidedly useful bird.



Common Kingfisher (*Alcedo ispida*).



THE PIED MYNA
(*Sturnopastor contra contra*).



THE PIED MYNA

(*Sturnopastor contra contra*).

ALTHOUGH almost as familiar a bird in North-Eastern India as the Common Myna, the Pied Myna has a much more restricted distribution, being found only east of a line drawn from Amballa to Hyderabad, in the Deccan, and Masulipatam, ranging to the extreme east of Assam. Where it occurs, it is common and conspicuous, both by its strongly contrasting colours of black and white and by its habit of hunting about over grass in the open, often in small flocks, and there is little chance of mistaking any other bird for this one. Our Plate gives a good idea of its coloration, so that there is no need to describe this here.

Like its commoner relative, the Pied Myna is a frequent attendant on cattle, picking up the various insects disturbed as the latter graze. It also hunts, almost always in small parties, over any open grassy spaces, catching and eating grasshoppers, crickets, caterpillars, ants and other insects and I have seen one extract and eat a large earthworm. Mr. C. W. Mason examined the stomach-contents of fourteen birds at Pusa and found that, of thirty-nine insects taken, one was beneficial, twenty-five injurious and thirteen of neutral value, and remarks that, on the whole, the Pied Myna is decidedly more vegetarian in its diet than the Common Myna. "When any *Ficus* fruit is ripe or a cereal crop, such as maize, the birds flock to these and at times will be found to feed on little else. They seem even more partial to *Ficus* fruit than the Common Myna. Some considerable damage is done at times to the various common cereal crops—maize, sorghum and paddy especially. Its



insect-food is much the same as that of the Common Myna, consisting as far as one can see in the field very largely of grasshoppers, small moths, etc." It may, however, be added, in justice to this bird, that nearly all of Mr. Mason's records were obtained in this case between February and June, when relatively little insect-food is available; I am inclined to think that a longer series of records, better spread over the year, would show that the Pied Myna is rather more largely insectivorous. Of course, like all similar birds with a mixed diet, it does levy toll on ripe grain, and the conspicuous colours of this bird rather tend to accentuate its presence in grain-fields, but against this must be placed the fact that it is continuously at work throughout the year in reducing the number of insects which, if unchecked, would undoubtedly do far more damage to the crops than is done by the birds. Further, it must be remembered that examination of the bird's stomachs only reveals their contents and does not indicate how the food has been acquired; in many cases, even when these birds are found feeding on cereal crops, the grain eaten is largely picked up off the ground and is almost always intermixed with a considerable proportion of insect-food. In the case of one bird examined at Nagpur, Mr. D'Abreu found it to contain four earwigs, two mole-crickets, two caterpillars, a bug, a batch of forty-seven insect eggs, and one red mite—which seems quite a useful effort for one meal; and it must be remembered that such a quantity is probably only a fifth or sixth of the total quantity eaten by one bird in the course of a single day. The occasional levy of a little grain, when it is abundant and when insect-food is scarce, as is usually the case during the harvest season, must not be begrudged to birds, such as Mynas, which on the whole are decidedly beneficial.

Cunningham, whose remarks on other birds we have frequently quoted in preceding articles, does not seem very kind to the Pied Myna and writes:—"They are not nearly such attractive birds as the Common Mynas; for their colouring is coarsely laid on in a




way that recalls that of certain of the ornithological inmates of a Noah's ark; their heads have a debased look, and they have neither the pleasant notes nor the alluringly familiar ways of their relatives. Like the latter, and very often in company with them, they spend their nights, save during the nesting season, in huge mobs, which, if possible, are even more vociferous than those of mynas. At sundown the din proceeding from such assemblies is often so overpowering as to render even the concerts of the crows or of the great autumnal crickets temporarily inaudible. Although roosting in and haunting gardens, they never show any desire to enter houses." In Bihar, this bird seems to be locally migratory; at all events, it seems to occur in large numbers at some times and at other times none are noticed during quite a considerable period.

The breeding season of the Pied Myna is from May to August but most birds lay in June and July. At the time of writing (4th June) there is a nest in a *Dalbergia* tree alongside my bungalow and this apparently contains young birds, as the parents are bringing in food, principally caterpillars apparently, but they are shy and it is not always easy to see what they do bring. The nest is situated in the fork, near the extremity of a bough, some twenty feet from the ground, and is a large untidy mass of twigs with a few strips of rags attached to the underside, relatively huge for the size of the bird, being about two feet in diameter, but by no means unduly conspicuous amongst the leafy foliage. This is quite a typical nest, these being built as a rule of straw, grass, twigs, roots and rags, with a deep cavity lined usually with quantities of feathers. Very rarely the nest may be placed in the cavity afforded by a hollow tree-trunk. Five eggs are laid as a rule, sometimes four, and occasionally as many as six. The egg is typically a moderately broad oval, considerably pointed towards one end, but sometimes pear-shaped, pale to sky-blue, sometimes tinged with green, unspotted and brilliantly glossy, in size about 28 millimetres long by 20 broad. The young, as in the case of most birds with

a mixed diet, are fed wholly on insects which are brought in by the parents.

The Pied Myna is protected under the Wild Birds' Protection Act, in the United Provinces, Bengal and Assam. Oates gives the vernacular names under which it is known as *Ablak maina* in Hindi, *Ablaka gosalik* and *Guialeggra* in Bengali, and *Venda gorinka* in Telugu.

A subspecies, occurring in the South of Assam and in Burma, has been separated under the name of the Burmese Pied Myna (*Sturnopastor contra superciliaris*) and another subspecies, which occurs in Siam and just enters our limits in Tenasserim, is the Siamese Pied Myna (*S. contra floweri*).



A young Babbler.



THE INDIAN PARADISE FLY-CATCHER
(*Terpsiphone paradisi paradisi*).



THE INDIAN PARADISE FLY-CATCHER

(*Terpsiphone paradisi paradisi*).

The paradise fly-catcher softly flits
Between the leaves with graceful flutt'ring tail
Of purest white: restless, he never sits
Upon a bough, but threads the leafy veil,
In dazzling contrast to the foliage dark,
The brilliant sunshine glinting on his plume;
His mate, a bird of sober brown I mark,
Sits peaceful near in her small nursery room.

Chenar Leaves.

THE Paradise Fly-catcher is probably one of the most striking of the birds that occur in India and is easily recognizable, at least in the case of the adult male, by the two extremely long tail-feathers which in some localities have earned their owner the name of Ribbon-bird.

The Fly-catchers form a group of birds feeding mainly on insects which are caught on the wing, the species of this group having very feeble feet, which incapacitates them from walking on the ground, their usual habit being to wait on some convenient branch and swoop down on their insect prey, which is taken in the air. Many other birds, such as the King Crow, also have this habit, so that we must not be taken to mean that every bird seen swooping down on an insect in the air is necessarily a Fly-catcher. This group of birds is characterized by the presence of numerous long hairs stretching from the forehead over the nostrils, these hairs lying horizontally and reaching in all cases beyond the nostrils and sometimes nearly to the end of the bill. Over fifty species of Fly-catcher occur within Indian limits and amongst these the



genus *Terpsiphone* is distinguishable by its crested head in conjunction with a tail as long as, or longer than, the wing. Three forms of *Terpsiphone* are recorded in the *Fauna* volume on Birds as distinct species but are now regarded as geographical races (sub-species) of *T. paradisi*; of these *T. paradisi paradisi* is recognizable by its long and pointed crest, reaching to the upper part of the back, whilst the crest is short and rounded, not reaching beyond the nape, in the case of the Burmese Paradise Fly-catcher (*T. paradisi affinis*), which occurs from Sikkim and North-East Assam to Tenasserim, and in the Nicobar Paradise Fly-catcher (*T. paradisi nicobarica*), which replaces the other two species in the Andamans and Nicobars.

The Indian Paradise Fly-catcher occurs throughout the whole of the Indian Region from Afghanistan and Kashmir to Ceylon, but in the North-East, eastwards of the Brahmaputra, it is replaced by the Burmese subspecies. It is found in most localities in the Plains throughout the year but appears to migrate locally to some extent, although there seem to be few exact records on this point; thus, Oates states that it seems to be everywhere a permanent resident, whilst Dewar says that it visits the Punjab in great numbers in summer for nesting purposes. We agree with Dewar that this bird does undergo a certain amount of local migration. In North Bihar, for example, it arrives about the end of March and leaves at the end of October, not being noticed during the cold weather. It is fairly common locally in Bihar but its distribution seems to be rather erratic: for example, I have never seen one at Pusa. In the United Provinces also it is a migrant, as Gill states that during the winter months this attractive bird occurs in most districts and is frequently met with in gardens and groves adjoining human dwellings, but the majority seem to migrate in summer to the Sub-Himalayan tracts, although a great many remain behind and breed sparingly throughout the Plains. So far as our experience goes, the Paradise Fly-catcher seems to occur most commonly in Northern India. I have seen it in considerable numbers around the



lake at Bhim Tal in Kumaon. In Calcutta it can hardly be called common, but stray specimens may be seen in gardens at all times of the year, and the same remark applies to Southern India.

The Paradise Fly-catcher when young is chestnut in colour, except for a black head and crest and whitish under-parts, and the female retains this coloration throughout her life. The cock bird, however, undergoes a very striking change of appearance, the two median tail-feathers growing to a length of sixteen inches (i.e., four times the length of the other tail-feathers) after the autumn moult of the second year, these long feathers being retained until the following summer, when they are cast. After the third autumn moult, these long feathers grow again and the plumage gradually begins to turn white (with the exception of the feathers of the head and crest, which remain black), the wings and tail being the first portions to be affected by the change; the adult male bird is thus partly chestnut and partly white for a time, and it is not until after the moult of the fourth autumn that the plumage of the body and tail becomes wholly white. Thereafter he remains white, with a black head and crest, and the two long white tail-feathers, looking like white ribbons, for the rest of his life, and is a most striking object when seen on the wing. As Cunningham says, "the first sight of one of them, floating softly along and seeming to swim through the air in a series of gentle impulses, gives rise to a very lasting mental impression. As one of the mature male birds flies along through the leafy coverts, in which they are most at home, the snowy whiteness of his long waving train gleams out in the light of the scattered sunbeams that struggle downward through the branches, and produces effects quite unlike those that attend the flight of any other kind of bird Even in their first dress, and before they have acquired their wonderful trains, they are strikingly beautiful. They have such full, bright, black eyes, such rich chestnut tints in the wings and tail, contrasting with the shining black of the head and the snowy white of their under-clothing, and their movements are so



exceptionally graceful that it is hard to cease from watching them, and when they are in all the glory of full dress, they must be to every one a source of wondering admiration as they leap lightly about from twig to twig and float hither and thither among the branches."

Oates writes that the notes of the birds of this genus are very harsh, but such a statement seems decidedly libellous and once again we cannot do better than quote Cunningham, who says that "whilst travelling about over the boughs, they continually utter twittering notes, with occasional louder calls, so like those of the Blue Fly-catcher that, until the birds come into view, it is impossible to make out which species one is listening to. Now and then, too, the male birds break out into sweet little songs. They are very lively and cheerful birds, always on the move; and the males constantly flirt their great trains about, separating and closing and undulating the long, trailing plumes in a wonderful way." The Indian name of *Shah Bulbul* also presumably indicates that this bird is by no means devoid of song. This bird does, however, at times (generally when on the wing) utter a decidedly harsh note.

All Fly-catchers are insectivorous as a general rule, although some species take fruits, berries and seeds occasionally. The Paradise Fly-catcher feeds on small beetles, flies, bugs, ants and spiders, as recorded by Messrs. Mason and D'Abreu from examination of actual stomach-contents. It is, therefore, together with the other species of this group, a useful bird to the agriculturist and its utility and beauty fully deserve the protection accorded to it by the Law in Bengal, Bombay and Burma (but not in other Provinces, apparently). In Mysore also it is presumably protected as being a bird of bright-coloured plumage.

The Paradise Fly-catcher lays its eggs in Ceylon and Travancore in February and March, in Southern India in March and April, and




in Northern India in May, June and July. The nest is usually a delicate little cup, never very deep and often quite shallow, composed, according to materials locally available, of moss, moss roots, vegetable fibres and fine grass, this last generally constituting the greater portion of the thin frame-work, which is bound around externally with cobwebs. The egg-cavity is from 2 to $2\frac{3}{4}$ inches in diameter and from one to 1.6 inches deep, and is often lined with horse-hair or fine grass. The nest is affixed to the thin branch of a tree, as a rule in a mango tree between seven and thirty (mostly between ten and twenty) feet from the ground. A favourite site for a nest is at the extreme end of a leafy, overhanging branch of a thick-foliaged tree, as in such a position the nest cannot be reached either from above or from below. Sometimes, however, the nest is placed low down, occasionally on the upper surface of a horizontal bough, in such cases usually at a place where there is given off a slender upward-growing twig which is built into the structure of the nest and thus serves to anchor it in position. Three or (less often) four eggs is the normal number, the egg being a rather long oval, somewhat pointed towards one end, about 20 mm. long by 15 mm. broad, in colour pale pinkish-white to warm salmon-pink, more or less thickly speckled, chiefly at the larger end (where there is a tendency to form an irregular cap) with rather bright, but somewhat brownish-red, spots, amongst which a few tiny, pale, inky purple blotches sometimes occur. Both the parents share in the duties of incubation of the eggs and feeding of the young, the male in such cases either having or not having attained his wholly white plumage; if he has done so, he is of course a very conspicuous object when sitting on the nest.

Our Plate gives a good idea of the male in wholly white plumage, as well as of the female and nest.

In the new edition of the *Fauna* volume on Birds, Stuart-Baker writes of the habits of this species as follows :—"There is no more

beautiful avian sight in India than a graceful, long-tailed, white, male Paradise Fly-catcher, as it flits backwards and forwards in the deep green shade of the Mango-orchards it so often haunts. They are tame confiding birds, frequenting gardens, open country and the vicinity of villages but they are also found in forest-land, especially deciduous light forests such as Sal, etc. They feed entirely on the wing, never descending to the ground and never searching the foliage for insects as some Fly-catchers do. Their ordinary flight is rather slow, the long tail undulating behind as the bird flies, but they are capable of very quick movement when hawking for insects."



Red-whiskered Bulbul.



THE INDIAN MAGPIE-ROBIN OR DAYAL
(*Copsychus saularis saularis*).



THE INDIAN MAGPIE-ROBIN OR DAYAL

(*Copsychus saularis saularis*).


... .. warbles in the throat
Of ev'ry bird, that hails the bloomy spring.
Or tells his love in many a liquid note
Whilst envious artists touch the rival string,
Till rocks and forests ring.

Hymn to Narayana.

PERSISTENT PESSIMISTS are wont to say that in India the most gorgeous flowers have no scent and that the birds have no song, but the Magpie-Robin, with many other birds, gives the lie to the latter part of this assertion, as the cock bird has a fine range of melody which is heard especially in the Spring. This is one of our most familiar birds occurring commonly, but never in large numbers, in every garden and cultivated area throughout the whole of Continental India and Burma and Ceylon, being especially a bird of the Plains, although it ascends the Himalayas to a height of about 6,000 feet. The cock is a perky-looking bird, rather smaller than a bulbul, glossy black with its lower parts sharply marked off in white, a white bar on the wing and the outer tail-feathers white; it runs along the ground after insects and has a habit of elevating its tail perpendicularly at the end of its run. The hen bird is marked much like her mate but shows greyish-brown in her plumage where his is black. The sexes pair for life and are commonly found in gardens, their habits being very robin-like. They are rather pugnacious birds, a pair generally keeping the whole of a garden to themselves, and the hens are, as Finn observes, of much more retiring habits than their mates. According to Hodgson, these birds

are caged for fighting purposes: "Fighting these tame birds is a favourite amusement with the rich (in Nepal), nor can any race of game cocks combat with more energy and resolution than do these birds."

The Magpie-Robin feeds on the ground, mostly on insects, its diet being varied by vegetable matter and an occasional earthworm. The late C. W. Mason investigated the stomach-contents of twelve birds at Pusa and found them to contain 142 insects, of which 7



Feathers of Magpie-Robin.

were classed as beneficial, 50 as injurious and the remaining 85 as neutral, the insects eaten including grasshoppers, crickets, ants, weevils, and other beetles, a few bees (perhaps not taken alive), wasps and cutworms. The injurious insects taken are of far more economic importance than the beneficial and this bird may certainly be reckoned as one of the gardener's friends. I have seen one tackle and kill a hawk-moth caterpillar which was almost as bulky as itself.




Breeding takes place from March to July, when a rough nest is constructed in any convenient hole in a tree or wall, but egg-laying takes place in the majority of cases during the second half of April and in May. Very occasionally the nest is placed in a bush, but almost invariably it is found in holes in trees, banks or walls or under the eaves of huts and sometimes in verandah roofs. Both parents assist in building the nest and in incubating the eggs. In the Plains the nest is usually composed of roots, grass, fibres and feathers, but in the Hills moss and lichens are largely used, the nest being saucer-shaped, broad and shallow, four or five inches in diameter with a central depression about an inch in depth; in any case, it is always put together very loosely and roughly. Five is the usual number of eggs laid, but we have seen as few as only two incubated ones, and the eggs are typically oval, neither very broad nor very narrow, averaging about 22 mm. long by 17 mm. broad, the ground-colour varying from greenish, greenish-white, to pale sea-green or delicate pale, only slightly greenish blue, but they are generally all streakily blotched and mottled with different shades of brownish-red, sometimes thinly and sometimes so extensively as to leave but little of the ground-colour visible, and in all cases the markings are most numerous at the larger end of the egg, on which they commonly form a conspicuous irregular mottled cap. We have taken one clutch of a pale greenish blue without a single mark on any of the eggs. The eggs in the same clutch vary very largely in the markings.

Our Plate gives a good idea of this bird, the cock bird on the right and the hen on the left, the butterfly represented being *Colias croceus feldi*, which occurs commonly all along the Himalayas, penetrating into the Plains in the cold weather.

In the south of Burma this bird is represented by a local race (subspecies), the Malay Magpie-Robin (*Copsychus saularis musicus*), in Ceylon and Travancore by the Ceylon Magpie-Robin (*C. saularis*



ceylonensis), and in the Andaman Islands by the Andaman Magpie-Robin (*C. saularis andamensis*).



Common Myna.



THE WEAVER-BIRD OR BAYA
(*Ploceus philippinus*).



THE WEAVER-BIRD OR BAYA

(*Ploceus philippinus*).

... the weaver-bird,
Chirped from her swinging nest ..

The Light of Asia.

THE Tailor-bird has already afforded us an example of admirable nest-building and the Weaver-bird has equal claims for admission to our circle of bird friends, which, so far as this series is concerned, must be strictly limited. We may, however, at once note a point of contrast between the two. The Tailor-bird makes every effort to escape its enemies by concealing its nest between leaves sewn together; the Weaver-bird, on the other hand, positively flaunts its nesting arrangements before our eyes and in many parts of India its nests—whole colonies of them, indeed—form conspicuous objects of the country-side. In some places scores of these nests may be seen, freely exposed to view, hanging from favourite trees, which are usually palms or *babuls* (*Acacia arabica*), trees which overhang water often being selected as it seems to be a *sine qua non* that the situation considered suitable for suspending a Weaver-bird's nest should have no other tree directly underneath it which might afford access to enemies from below. For a nest swinging up aloft on the tip of a long slender palm-shaft is singularly inaccessible to most would-be marauders.

Although the nests are familiar enough, the bird which constructs them is less so. The Weaver-bird belongs to the large family of the Finches and at most times of the year looks very like a hen House-sparrow, being a small, thick-billed, reddish-brown bird. Towards the beginning of the Rains, however, the cock bird dons



his breeding plumage, his head, neck and breast becoming a beautiful golden-yellow and his chin turning almost black.

With the onset of the Rains, nest-building commences, either a new nest being built or an old one, of the previous season, patched up and put into good repair. The process of patching up old nests, which are easily distinguishable by the difference in the colour of the grass, and building new ones can often be seen going on in the same tree. The nest itself is strongly woven with strips of leaves of grasses, plantain or palms, strips from leaves of wild species of *Saccharum* being used most commonly. These strips are prepared by the bird itself, which seizes a leaf in its beak and makes a notch at the edge near the base of the leaf; it then grips with its beak the edge of the leaf above the notch and jerks its head away so as to tear off a strip along the edge of the leaf; by flying off with the end of this strip in its beak, the strip is usually pulled off along the length of the leaf; but sometimes, at the first effort, if the leaf is tough, the bird is pulled backwards and hangs suspended by the strip in its beak, so that several attempts have to be made to detach the strip required for nest building. Sometimes the bird bites a second notch on the edge above the first and at a distance allowing for the length of strip which it considers necessary. The strips thus collected are wound securely around the branch or leaf from which the nest is to be suspended and, as more strips are brought in, these are added, securely wound and plaited together, until there is formed a long stalk from which the nest proper is suspended. This stalk is usually about four or five inches long but may occasionally extend to as much as a foot. Having completed the suspensory stalk, the birds then expand its lower end into a bulb-shaped structure, which is usually about five and a half inches in diameter in one direction and four inches in the other. At this time, having determined where the egg-chamber is to be, the birds construct a strong transverse bar or loop, a little to one side of the centre of the chamber, this bar forming a division between the egg-chamber and the long tubular



entrance. At this stage of construction the nest resembles an upturned basket, the loop representing the handle. Up to this point in the construction of the nest, both sexes have done the same kind of work in collecting fibre-strips and weaving these into the nest, but, when this loop has been completed, the female bird takes up her position on it, leaving the cock henceforth to procure more material for building and to work from the outside of the nest, whilst she works from the inside, both of them pushing and drawing in the fibres through the walls of the nest so that everything is plaited together smoothly. The little builders seem to enjoy themselves thoroughly, the cock bird especially being industrious, emitting a cry of delight each time that he brings in a beak-full of fibre and often bursting into song during the process of weaving material into the nest. The egg-chamber is now finished on one side of the loop and on the other side the walls of the nest are prolonged downwards into a long tubular entrance, about two inches in diameter internally, and usually about six inches long, but occasionally twice this length. The male bird often continues building on to this tubular entrance even after the eggs have been laid and are being incubated by the female. The lower end of the tubular entrance is loosely woven so as not to afford any firm support to enemies attempting to plunder the nest. The birds themselves when entering the nest close their wings and shoot perpendicularly upwards through the tube; it is marvellous how they can do this without running their heads through the top of the egg-chamber or even apparently shaking the nest. This tubular passage is used as an entrance to the nest by the parents whilst nest-building and incubation are proceeding; but, when the eggs have hatched, the food brought to the nestlings is passed in to them through small holes pierced through the sides of the egg-chamber. The presence of such holes is a sure sign that the eggs have hatched.

It will now be apparent what a hard nut has to be cracked by any would-be plunderer of the nest, which is placed high up



out of reach of any non-climbing animals. Even a good climber, such as a squirrel, rat, snake or lizard, will find little to cling to on the tip of a palm-leaf and, arrived there, has to negotiate a distance of at least eighteen inches to reach the entrance to the nest, whilst the lower portion of the entrance-tube is too flimsily constructed to yield any foothold. Plainly visible, its inaccessibility is its sure defence and it can well defy most marauders.

There is still one point to be mentioned about these nests and that concerns the lumps of clay which are stuck onto them in odd places. Jerdon notes that he found in one nest about three ounces of clay in six different places. Many theories have been advanced in explanation, a very popular idea in India being that the bird uses these clay patches as *points d'appui* on which to stick glow-worms to illuminate the interior of its nest. A more probable explanation is that the clay is applied to balance the nest more correctly, to prevent it being blown about by every gust of wind and to keep it steady whilst the birds are entering and leaving it.

Two is the normal number of eggs laid but occasionally three or four are found. As many as ten have been noted, but, in cases where there are so many, they are probably the product of more than one bird. The eggs are usually found in July and August. They are pure white in colour, without any gloss, typically rather long ovals considerably pointed towards the smaller end, and measure about 20 by 15 millimetres.

The Baya is found throughout India and is now divided into four subspecies, which are the Baya (*Ploceus philippinus philippinus*) found in Ceylon and the greater part of India; Finn's Baya (*P. philippinus megarhynchus*), a local race found in the Himalayas about Nainital; the Eastern Baya (*P. philippinus passerinus*), found in the lower Himalayas and Hills in Bengal, Assam, North Burma and Siam; and the Malayan Baya (*P. philippinus infortunatus*)




which occurs in the Malay Peninsula and Siam, only entering our limits in Tenasserim. It is the typical race which is shown on our Plate. Besides the Baya, three other species of Weaver-birds are found in India, of which the Indian Striated Weaver-bird (*Ploceus manyar flaviceps*) has the feathers of the breast streaked longitudinally with black; this bird is not uncommon in localities in Northern India providing suitable rushy, reedy cover, in which the nests are placed, the nests being much like those of the Baya but without the long pensile support.

Like most of its relatives, the Weaver-bird is largely graminivorous, feeding on seeds of grass, paddy, millets, and weeds, but a certain proportion of insect food, mostly small beetles and caterpillars, is taken. It cannot be claimed as a useful bird, and during times of forest fires the nests sometimes burn through at the base and may then be blown, all ablaze, for hundreds of yards into areas which would otherwise escape from the fire.

Its feeding habits make it comparatively easy to keep the Baya as a cage-bird and it is often so kept, young birds being offered for sale in Calcutta during August. Given a large enough space, the Baya will weave its wonderful nest in confinement, but requires all the space for itself for, as Cunningham says, "they are very undesirable additions to any aviary containing other kinds of small birds, as they are very aggressive, and are possessed by a deeply-rooted desire to hammer in the skulls of their neighbours, which, as Abdur Rahman in his autobiography remarks of a Baluchi tribe of similar disposition, 'naturally causes disputes'".

Young Bayas are readily tamed and easily acquire tricks, such as threading beads, drawing up little buckets of water or of seed, or loading and firing off a toy cannon. Lockwood Kipling tells of one which flew up to a tree at the word of command, selected a flower or leaf, plucked it, and, returning, placed it daintily between

its master's lips. There is no doubt that it is an intelligent bird and it is therefore a favourite cage-bird. In the Punjab a popular proverbial rhyme contrasts the house-building talents of the Weaver-bird with the helplessness of the shelterless monkey which cannot protect itself against the weather in spite of possessing human hands and feet. "This verse," says our informant, "is often quoted for the benefit of idle boys and girls who object to learn," much in the same way as the little busy bee is held up for infantile admiration in Western lands.



Feathers from tail and wing of female Koel.




PLATE XIV.

THE INDIAN HOUSE-SPARROW
(*Passer domesticus indicus*).



THE INDIAN HOUSE-SPARROW

(*Passer domesticus indicus*).

A little cock sparrow sat on a tree,
Looking as happy, as happy could be,
Till a boy came by with his bow and arrow :
Says he, "I will shoot the little cock sparrow.
His body will make me a nice little stew,
And perhaps there'll be some for a little pie too,"
The cock sparrow said, "I'll be shot if I stay,"
So he stretched out his wings and flew right away.

Old Nursery Rhyme.

THE Indian House-Sparrow needs little introduction to our readers, as it is only too familiar throughout India, occurring abundantly in all towns and by no means uncommonly in all country districts. The Indian race differs slightly from the form so familiar in Europe in being usually much whiter about the sides of the head and in having more black below the eye and at the base of the cheeks, so that the Eastern form looks more brightly coloured than the Western, but these characters vary considerably and the Indian race is not now considered as more than a local form of the European species. However, as Hume says, "What is in a name? Call him *domesticus* or *indicus*, it doesn't alter his depraved nature, does not make him one whit less detestable—only there is a certain *lucus a non lucendo* sarcasm involved in the Linnean name that aggravates.

"If domesticity consists in sitting upon the punkah-ropes all day, chit, chit, chit, chattering ceaselessly when a fellow wants to work, banging down in angry conflict with another wretch on



to the table, upsetting the ink, and playing old Harry with everything, strewing one's drawing-room daily with straw, feathers, rags, and every conceivable kind of rubbish in insane attempts to build a nest where no nest can be—if, I say, these and fifty similar atrocities constitute domesticity, heaven defend us from this greatly lauded virtue, and let us cease to preach to our sons the merits of *domestic* wives! Conceive a wife evincing similar tendencies! Why, there isn't a jury in the country who would not return a verdict of 'sarve her right,' even if the unhappy husband should have wrung her neck before the golden honeymoon had run out."

The above condemnation of the Sparrow is certainly sweeping but there is no doubt that this bird must be regarded as a decided pest of all well-ordered households in a country such as India, where the numerous open doors and windows afford easy entrance and exit and the lofty rooms and verandahs, with their heavy projecting cornices and numerous chinks and crevices, provide the Sparrow with such a convenient variety of suitable nesting-places. Noisy, pugnacious and untidy will perhaps sum up the character of this bird. Cunning, crafty, hardy, and well-nigh omnivorous, it has become a parasite of the human race and is well described by the epithet of "the avian rat," flourishing especially in localities such as towns, where human activities have upset the normal balance of Nature and destroyed or driven away the natural checks, especially the birds of prey which help to reduce the numbers of this bird in rural areas.

In Europe, America and Australia the House-Sparrow is justly regarded as a very serious pest of growing crops and, although relentless persecution is carried on and its numbers kept in check, it is still able to inflict annually loss and damage that can only be reckoned in millions of pounds. In India, things are not quite so bad mainly because the Sparrow is kept in check by natural



causes except in large towns. Not only do its disorderly and noisy habits make it an active nuisance in and around houses, but its food consists largely of vegetable matter acquired at the expense of its rightful human owners. Mr. Mason examined the stomach-contents of eight birds at Pusa and Mr. D'Abreu of eight others at Nagpur and in all cases they were found to contain seeds of various grasses, including cultivated forms such as rice and oats; none of the birds from Nagpur contained any insects and of the eight from Pusa only two contained, between them, two small weevils and another small beetle. The young nestlings, it is true, are fed largely on an insect diet, composed chiefly of caterpillars, but as the young grow older the proportion of insect food given them is diminished until, when they are about three weeks old, their diet is made up almost wholly of grain. The same story is revealed in Australia, whither this bird has been introduced with the unfortunate results so often encountered in the case of introduced animals and plants, and in a Bulletin on "The Food of Australian Birds" we read that, of 127 Sparrows examined, sixty-four birds contained wheat and maize seeds and it is remarked that this bird is "a pest anywhere, in spite of the fact that it eats many insects." Besides grass-seeds and occasional insects, the Sparrow does not disdain the smaller fruits and in the mulberry season an extra annoyance is added to its presence in houses owing to its partiality for this fruit.


The nesting season is principally from February to May but two or more broods are undertaken annually and breeding continues throughout the year, the semi-domesticated conditions of this bird's existence not restricting it to any one season of the year. Its nest is a shapeless bundle of straw, grass, rags, wool or anything else obtainable, thickly lined with feathers and stuffed into any available hole or recess in or about houses, walls, old wells, etc., or rarely even in the centre of a thick bush. If a tree or a climber on a wall be chosen for the nesting site the nest is better



made and is often a substantial dome-like structure with the entrance at the side, but its position is usually betrayed by long untidy pieces of straw left trailing outside. Five or six eggs, sometimes even more, are laid at each breeding season, the eggs being typically somewhat elongated ovals and but little pointed, either greenish, greyish, or yellowish-white, marked with close frecklings, fine lines or smudgy streaks of dull dingy sepia, olive, yellowish or purplish-brown, these markings being sometimes sharply defined, and often showing a tendency to form a blotchy, mottled, ill-defined cap at the larger end. The eggs vary much in size but average about 20 mm. long by 15 mm. broad.

Besides the natural enemies of the older birds, the nestlings are attacked whilst in the nest by the grubs of a fly, *Passeromyia heterochæta*, which lays its eggs in the nest and whose larva buries its anterior extremity in the skin of the nestling, usually under the wings or the legs, and sucks its blood. This fly is widely distributed in Africa and has been found in China and in India at Pusa and Coonoor and is probably widely distributed in India. The numbers of the adult birds are largely kept down in the *mofussil* by hawks and probably also by the Indian Vampire Bat (*Lyroderma lyra*) which hunts by night along the hedges in which these birds congregate in the evening to sleep in flocks, this habit affording an opportunity of netting them in quantity when it is desired to reduce their numbers.

The Indian House-Sparrow occurs throughout the entire Indian Empire and in Ceylon but is not found in the Andamans or Nicobars or in the south of Burma. It ascends the Himalayas to moderate elevations. It is only too abundant in most parts of its range but is rare in some localities. In Sind there is a local race, the Rufous-backed Sparrow (*Passer domesticus pyrrhonotus*).



THE WHITE WAGTAIL
(*Motacilla alba*).




THE WHITE WAGTAIL

(*Motacilla alba*).

INDIA has many birds which visit the Plains as winter tourists only, passing the summer months either in the Hills or in "the back of beyond" far to the north of the mighty Himalayas. Everyone is familiar with the fact that ducks, for example, migrate in this way but it may be news to some of our readers that most of our common wagtails also act in this manner, visiting India only in the cold weather and passing the summer in Northern Asia. About a dozen different forms are found in India but only one (*Motacilla alba maderaspatensis*), a black wagtail with a conspicuous white eyebrow, looking not unlike a magpie-robin but never carrying the tail erected, is a permanent resident. All are very similar in general build, being slenderly built, dainty-looking birds, mostly coloured in mixtures of black, white and grey, occasionally yellowish or greenish, their delicate slimness harmonizing well with their lightness of gait as they run with great speed after their quarry, never hopping but sometimes making little sallies into the air, and constantly wagging their tails whose outer feathers are always white and conspicuous during flight; whence their popular name. Then, as Cunningham has well remarked, they are so alluringly tame, merely running on in front of one and expostulating at being disturbed; and, if persistently followed along a narrow path, making off on a brief, undulating flight to pitch anew a little way ahead in a way that gives one the fullest opportunity of becoming familiar with them. They are so tame that they often come into verandahs of bungalows and pick up insects off the floor. The migrant species

usually arrive in India during September and remain with us until the end of the cold weather, the White Wagtail leaving Bihar before the end of April.

Wagtails feed mostly on small insects, chiefly flies, small beetles, ants and caterpillars, but occasionally pick up seeds also. The insects eaten are not of any beneficial kinds and in many cases are injurious to crops, and therefore these birds as a whole must be



White Wagtail.

reckoned amongst the farmer's friends. It is a pity that most of them are absent from India during the Rains, when insect life is so abundant. The particular species which we have selected as a representative of this group is the White Wagtail (*Motacilla alba*), which Dewar briefly describes as, general colour of plumage grey; face, chin, and throat white, back of head and nape black; a black patch on the breast, the remainder of lower plumage white; the wings white with much black on them; the middle tail-feathers




black, the outer ones white. Our Plate gives a good general idea of its appearance. It has a very pleasant note. It is found throughout most parts of India during the winter months but does not occur in Southern India nor South of Moulmein in Burma. The Masked Wagtail (*M. alba personata*) is by some authors regarded as a distinct species, but by others as a form of *M. alba*, from which it differs by the ear-coverts and sides of neck being always black, whereas in *M. alba* these parts are always whitewashed. The distribution of *M. personata* much resembles that of *M. alba*, but it does not occur east of Calcutta and is a constant resident, and breeds, in North-West India.

The White Wagtail is a regular visitor to England during its time of migration from the Continent of Europe and breeds in England at times and has been known to pair there with the Pied Wagtail (*M. lugubris*), a species which does not occur in India. Its nest has been found there in such odd places as in a Sand Martin's burrow, in the middle of a strawberry bed, and amongst a Virginian creeper growing over trellis-work. The eggs are bluish-white, speckled with black. So far as we know, it never breeds in India, and Hume in his *Nests and Eggs of Indian Birds* omits the White Wagtail altogether but mentions the Masked Wagtail, *M. alba personata*, which is perhaps a subspecies of the White Wagtail, as breeding in Afghanistan in May and June, the nest being usually placed in holes, under large stones, in or near beds of rivers.

The habits of the White Wagtail resemble those of the Pied. Dr. A. G. Butler gives some interesting notes on the latter species, in the course of which he says, "their power of turning in the air is astounding; few insects, however eccentric their flight, can hope to escape them. If a Wagtail is on the ground and it sees an insect flying towards it, instead of starting madly forward to meet its prey, it excitedly watches all the insect's movements and suddenly (when the latter is almost overhead) the agile bird rises with a

rapid spiral movement, which looks almost like a somersault, the snap of its mandibles is heard, and all is over." He also writes, "even when caught wild, most examples of *Motacilla* soon become tame if kindly treated; they are easy to feed, living for years upon crumbled household bread, yolk of egg and ants' cocoons, moistened (either by the addition of a little water or mashed potato) and a few insects, their larvæ or spiders from time to time."

As most species of Wagtails are not permanent residents and do not breed in India, they have not received legislative protection in most Provinces, except in Bengal where they are protected throughout the year and in Burma where they are protected in reserved forests from being hunted, shot, snared or trapped for purposes of trade.



Young Babbler.



THE INDIAN TREE-PIPI'T
(*Anthus hodgsoni*).



THE INDIAN TREE-PIPIT.

(*Anthus hodgsoni*).

With love exceeding a simple love of the things
That glide in grasses and rubble of woody wreck ;
Or change their perch on a beat of quivering wings
From branch to branch, only restful to pipe and peck.

Meredith—*Melampus*.

THE Pipits, contained in the genus *Anthus*, are all rather dull-coloured birds in shape like Wagtails, to which indeed they are closely allied, and may be recognized by their streaked upper plumage and comparatively short tail. About a dozen different species occur within the Indian Empire and resemble one another very closely, so that they are difficult to distinguish in the living condition, their identification depending on such small points as the length of the hind claw. To the ordinary observer, then, all the Pipits look much alike and a bird which is seen to look much like the one depicted in our Plate may be set down as a Pipit but may not necessarily be the Indian Tree-Pipit. Pipits as a class are very highly parasitized by the Common Cuckoo, several species of these birds having been found to be victimized in this way, but apparently there is no record of the subject of our present article having been made use of as a foster parent.

The Indian Tree-Pipit is another bird which, like the White Wagtail, is only a winter visitor to the Plains of the greater portion of the Empire, being found as far West as the Sutlej Valley in the Himalaya and Rajputana and Gujarat in the Plains, and as far South as the Palni Hills. In Bihar and Bengal it appears



amongst the very earliest of the autumnal immigrants and remains in considerable numbers until about April, but during the summer it retreats to the upper ranges of the Himalayas and beyond to Siberia, North China and Japan. When they first arrive in India in the autumn their plumage is in fine condition and brightly marked but their colour becomes decidedly duller during their stay.

Tree-Pipits are very partial to mango groves and are rather shade-loving birds. However, they also frequent open spaces, such as garden paths, much in the same way as the Wagtails, and similarly keep their tails in constant rocking motion whilst pacing over the ground. When preparing to take flight they have a curious habit of swaying themselves about for a time, and when alarmed they fly up to the nearest tree and walk about on the branches in a decidedly un-wagtail manner. They are fairly social and frequently go about in small flocks, and feed on the ground on seeds of grass and weeds and on small insects. Seeds are oftener consumed by these birds than by the Wagtails, which are practically totally insectivorous. The late C. W. Mason examined the stomach-contents of sixty-seven birds at Pusa and Mr. D'Abreu those of three birds at Nagpur, and in all these seventy cases the birds contained seeds of weeds, injurious or neutral insects, and a few small snails. From an agricultural point of view, therefore, the Tree-Pipit may be put down as a decidedly beneficial bird, in spite of which fact it is commonly sold in the markets as an "Ortolan."

The Tree-Pipit does not breed in the Plains and indeed nests but sparingly within our limits and then only at an elevation of eleven or twelve thousand feet where it has been found nesting in Northern Kumaon. Here it affects by preference the more open grassy slopes in the immediate vicinity of woods; these open glades are thinly covered with trees and overgrown with beautiful



thick, soft, velvety grass, about a foot high, with occasional tussocks, under which are concealed the nests, built of dry grass blades, or of green moss lined internally with fine grass-stems. The eggs, which are very large for the size of the bird, are shortly described as "very black-looking."

Mr. C. M. Inglis found the nests of this bird at Sandakphu (elevation 12,000 feet) in the Darjiling district on 2nd July 1904 and also at Tonglu (elevation 10,000 feet) on the following day. Sandakphu has a kind of alpine pasture, the ground being carpeted with many beautiful herbaceous plants of alpine character, such as Primulas of several kinds and Aconite; also amongst the rocks occurs a common stiff-branched shrub (*Potentilla fruticosa*, Linn.) with handsome yellow flowers and silvery foliage. The nests were found on the sloping grassy banks, generally hidden under a clump of grass, often also close to the rocks. On 2nd July young birds had already hatched out but fresh eggs were obtained on the following day.

During the 1921 Mount Everest Expedition a specimen of the Indian Tree-Pipit was collected on East Everest at a height of 17,500 feet and this species was observed migrating on the side of Mount Everest at an elevation of 20,000 feet.

The name *maculatus*, under which this bird is described in the "Fauna" volume, must be abandoned as preoccupied and its correct name is apparently *Anthus hodgsoni*.

The Indian Pipit (*Anthus richardi rufulus*), or Indian Tit-Lark, is a very similar species which is likely to be found in any garden in which the grass is allowed to grow long. Unlike the Tree-Pipit, it remains with us throughout the year, breeding in the Plains and Lower Hills of India and Burma. Where suitable open grassy spaces are available, this bird builds at the beginning of the hot

weather, its nest being placed in the shelter of tufts of grass or under clods or similar objects affording some shelter. The nest is composed of a thin lining of dried grass placed in a slight depression in the ground, which must often get flooded out during heavy rain, but the occasional destruction of the first brood in this way is compensated by the rearing of a second brood about the end of May.



Hen Koel.

The female Koel, with her brown white-barred plumage (see figures of feathers on page 74, *ante*) is very different in appearance from the male Koel, which looks like a larger edition of the Drongo (see Plate VI). The manner in which the two sexes combine to parasitize the Indian House-Crow is described on page 13.



THE INDIAN WHITE-EYE
(*Zosterops palpebrosa palpebrosa*).



THE INDIAN WHITE-EYE

(*Zosterops palpebrosa palpebrosa*).

In his tam parvis atque tam nullis, quæ ratio,
Quanta vis, quam inextricabilis perfectio.

Pliny.

ALL of the birds with which we have dealt in our previous papers have been not only common, but also conspicuous species probably well-known to all who live in India. The subject of our present note, however, although sufficiently common in all localities in the Plains, is decidedly not a conspicuous bird and is easily passed over unnoticed.

The White-eye is a small bird, two-thirds only of the size of the common sparrow, in colour greenish-golden yellow, greyish-white below, with a bright yellow chin and throat and a patch of yellow beneath the tail; around the eye is a ring of white feathers, whence this bird derives its popular English name; on this account it is also sometimes called the Spectacle Bird. It seems to have no vernacular Indian name. This white eye-ring is distinctive and readily permits the identification of its owner as a member of this group, which was classed in the *Fauna* volume on Birds in a sub-family connecting the Babblers and the Thrushes but which is now placed as a separate family, the Zosteropidæ. Five forms of White-eye are recognized in the *Fauna* volume on Birds, but it seems probable that some of these are merely geographical races of the common Indian species. In Mr. Stuart-Baker's more recent Hand-list he recognizes only four species, of which *Z. palpebrosa* is split up into six subspecies.



All the White-eyes are small birds, which go about in flocks, frequenting trees and bushes, whose leaves they search for small insects, varying their food at times with small buds, seeds and fruits. Like most other birds which associate in small parties, they keep up a constant call one to another, their call being best described as a cheeping twitter.

The food of the White-eye consists of a mixed diet of insects and vegetable matter. Of sixteen birds examined at Pusa by the late C. W. Mason, eleven contained small buds, seeds and wild fig fruits, and five contained insect food, consisting mostly of weevils and ants. One bird, examined by Mr. D'Abreu 'at Nagpur, had been feeding on berries of *Zizyphus cenoplia*. The White-eye has also been recorded as damaging ripe mangoes and guavas, and will eat plantains when in captivity. In spite of a decided taste for ripe fruit, however, this bird cannot be called a pest of fruit-trees and probably does a considerable amount of good by picking off small insects throughout the year.

The White-eye occurs throughout the whole of India, both in the Plains and in the Hills, in Ceylon and the Nicobars, and in Burma. It breeds, according to locality, from January to September, but the beginning of the hot weather is its normal breeding season in most localities, and at that time the cock bird sings a sweet little song. In North Bihar it breeds from April to July, the earliest nest having been found on 1st April and the latest on 23rd July. With the exception of one nest found in a jak tree all the nests found have been in mango trees. Sometimes there are two broods in the year. The nest is suspended between two twigs or more rarely in a fork of a branch, at any elevation from less than a foot to sixty feet from the ground, but the majority build at comparatively low elevations, between two and six feet from the ground. It varies much in size and in the materials of which it is composed but is always a soft, delicate little cup, sometimes




quite shallow, sometimes decidedly deep, and almost always is suspended like a miniature Oriole's nest, from small twigs or leaves. The parent birds set to work with cobwebs and first tie together two or three leafy twigs, to which they intend to attach their nest, and then use any available fine fibrous material (such as vegetable fibres, fine roots, thread, floss silk or cobwebs) to complete the outer fabric of their very beautiful and compact nest. As the work progresses, more fibres and cobwebs are used, so that the nest might be taken by a casual observer for an accidental accumulation of rubbish caught in a spiders-web. The interior is lined with silky seed-down, hair, feathers, moss, fine grass, etc., according to the materials available. The external diameter of the nest is about $2\frac{1}{2}$ to 3 inches and the depth varies from one to two inches, the egg-cavity being about $1\frac{1}{2}$ inches in diameter. Two eggs are laid as a rule, occasionally three and very rarely four. The egg is rather long and pointed at the smaller end, about 16 mm. long by 12 mm. broad, in colour uniformly very pale blue or greenish blue, without any markings; but occasionally the egg is capped with a zone of rather deeper and purer blue.

Our Plate gives an excellent idea of this bird and of its nest as built in the fork of a branch of a mango tree.

An interesting account of the feeding of the young is given by B. B. Osmaston, who says that he observed one of the parent birds place rather a large insect into the wide open mouth of one of the young which in spite of many efforts appeared to be unable to swallow it; the other parent, which was watching operations from above, seemed to grasp the situation at once, for he (or she) bent down and put a small drop of liquid into the widely gaping mouth; the process of giving small drops of liquid was repeated in front of the observer quite four or five times, and he was so close that there could be no mistake; eventually the insect became soft and more manageable and the little bird's efforts were crowned with success.

The Indian White-eye is commonly seen in the Calcutta bird market at Tiretta Bazaar and is easily kept in confinement on a diet of bread and milk, soft fruit and small insects, and is well worth keeping.



The Dayal.



THE PURPLE SUN-BIRD OR HONEY-SUCKER
(*Cyrtostomus asiaticus asiaticus*).



THE PURPLE SUN-BIRD OR HONEY-SUCKER

(*Cyrtostomus asiaticus asiaticus*).

. In the mango-sprays
The Sun-birds flashed.

The yellow Sun-birds whirled from bloom to bloom.

The twitter of the Sun-birds starting forth
To find the honey ere the bees be out.

The grey eggs in the golden Sun-bird's nest
Its treasures are

The Light of Asia.

THE Magpie-Robin described in a previous article is by no means the only bird whose cheery notes enliven our compounds. Conspicuous amongst the smaller feathered songsters and appealing both to eye and ear are the exquisite little Sun-birds or Honey-suckers, of which the males, clothed in metallic colours, are amongst the most beautiful of the birds which occur in India. We have none of those lovely little gems, the Humming-birds, which flash splendidly from flower to flower in South America, but we may almost imagine that we have when we see a tiny bird, clad in the most brilliant purple plumage, hovering over a flower with rapidly vibrant wings and extracting nectar with its long tongue, quite in the manner of a Humming-bird or Hawk-moth. Such a bird is almost sure to be the male of one of our species of Sun-birds belonging to the genus *Cyrtostomus* (*Arachnechthra*), of which about eight species occur within Indian limits, but of these only three are commonly distributed in the Plains: Loten's Sun-bird (*Cyrtostomus lotenius*), which occurs in Ceylon and Southern India, being wholly dark



metallic purple above, with a snuff-brown abdomen and a long curved beak; the Purple Sun-bird (*C. asiaticus*), which is found throughout the Plains of India and in Ceylon, but is only a summer visitor to the Punjab, being very similar to the preceding, but with a violet black abdomen and a shorter curved beak; and the Purple-rumped Sun-bird (*C. zeylonicus*) which occurs in Ceylon, Southern and Central India, but not in Northern India or Burma, and is distinguished from the other two by its bright yellow lower plumage and crimson back. In Calcutta the Purple-rumped Sun-bird is the commonest species and occurs in all gardens; the male is a very beautiful little bird, bright canary-yellow underneath with rich reddish-brown wings and back, a lovely violet throat and a gleaming crown of metallic green on its head. The females of these three species do not possess the brilliant metallic coloration of their mates and have to be content with a dress which is earthy-brown or greenish-brown above and yellow beneath.

The above-mentioned birds belong, as already stated, to the genus *Cyrtostomus*, perhaps better known under the old name of *Arachnechthra*, but we have altogether some twenty-five species of Sun-birds within Indian limits and these are divided into four genera, the largest genus in point of numbers being *Æthopyga*, containing those birds in which the males have lengthened middle tail-feathers and yellow rumps. In this genus the commonest Plains species is the Himalayan Yellow-backed Sun-bird (*Æthopyga siparaja seherice*), a bird with most of the upper plumage crimson, as is likewise the lower plumage as far as the breast; it also has a purple moustache-like streak and the lengthened tail-feathers are metallic green. This Sun-bird is common in the Plains of North-Eastern Bengal, Assam and Cachar, and is also found along the foot of the Himalayas, which it ascends to a considerable height. It is parasitized in Assam by the Emerald Cuckoo (*Chrysococcyx maculatus*), and Mr. Inglis has taken several of this Cuckoo's eggs from the nest of this Sun-bird.



In their habits all the species are very similar, frequenting flowering trees and shrubs, extracting the nectar from the flowers with their long tubular tongues, either occasionally hovering on the wing or more frequently clinging to slender twigs. But, besides carrying out the poet's dictum,

“ For he on honey-dew hath fed,”

these little birds variegate their diet with small insects, for catching which their long bills are admirably adapted, both mandibles being serrated along the terminal third of their length. Small spiders, caterpillars, beetles, bugs and flies, probably in most cases themselves visitors to flowers, fall a prey to these birds. We have also seen the Purple Sun-bird picking insects from off the ground and also flying up and catching them on the wing.

Besides being useful in helping to reduce the numerous insects which haunt our gardens, Sun-birds are also directly beneficial by helping to pollinate many flowers. Writing of *C. zeylonicus* in Calcutta, Cunningham says that “ the curious narrow tubular flowers of *Hamelia patens* are very special favourites, owing to the large store of nectar in their lower ends ; and during the whole time that the shrubs are in flower they are sure to be alive with honey-suckers every morning. In this, and doubtless in many other cases, they seem to play a very important part in securing cross-fertilisation ; for, by the help of a field-glass, one can clearly see that every time their bills are withdrawn from one tube and thrust into another, they are thickly smeared with golden pollen ; and when flowers from which they have just been feeding are examined, the long oval stigmas will be found coated with adhering grains. In rifling the flowers, therefore, they confer a benefit on the plant, and do not play the part of mere robbers, like the great brown hornets, who share their liking for the nectar, but who, in order to



reach it, drill holes through the corollas below the level at which the anthers lie.

“Curiously enough, they do not seem to care for the fluid in the corollas of the silk-cotton trees, which is so attractive to so many other kinds of birds that the trees, when in full bloom, become noisy and riotous taverns thronged with excited toppers. The unopened flowers of *Hibiscus rosa-sinensis* are greatly frequented in the early morning on account of some attractive material to be found at the bases of the petals. *Erythrinas* are also very popular; the clusters of their bright red flowers are very often alive with a throng of clinging and fluttering little thieves; and an even more charming picture presents itself when the latter are busy among the deep green foliage and tufted crimson inflorescence of *Hæmatocephala Hodgsoni*.”

In his volume, *Some Indian Friends and Acquaintances*, Cunningham gives a long and charmingly written account of the habits of *C. zeylonicus* in Calcutta, from which we would willingly quote at further length, did space permit, and we can only advise our readers to refer to Chapter XI of his book.

The Purple-rumped Sun-bird is the dominant species in Calcutta, but in Bihar our common species is the Purple Sun-bird, which is depicted in our Plate. This is a less sociable bird, more often seen alone and not in company with others, as one sees *C. zeylonicus*, but it is a much better songster than the latter, singing much like a canary; during the Rains, however, in the non-breeding season, it seems to become more sociable and a dozen or more of these delightful little birds may often be seen feeding on one flowering bush, fluttering and hopping about or swinging head downwards suspended by their strong prehensile feet whilst they imbibe the nectar from the flowers, the still, damp air alive with their cheerful twitterings. The cock bird in the breeding season is a truly gorgeous



little creature, appearing a blue-black at a little distance ; but when seen at close quarters the colour is a metallic violet-blue or greenish, the colour changing according to the intensity and angle of the light on it so that it may appear shining blackish-purple or green or more often mauve.

".....cecu nubibus arcus
Mille iacit varios adverso sole colores."

Under the base of his wings he carries a large tuft of mixed orange-red and yellow feathers, which is ordinarily concealed under the wings, but which projects when the bird is settled for repose and is displayed when courting. Like human bipeds, however, the Purple Sun-bird does not wear his wedding garments throughout the year, for after the nesting season is over he doffs them and assumes female plumage, retaining only a dark metallic violet streak from chin to abdomen as a mark of his sex. It may be added that there is a certain amount of difference of opinion as to whether the purple plumage of the male is or is not retained throughout the year, Messrs. Oates and Dewar contending that, when once assumed, it never changes ; but Finn, in an article in the *Journal of the Asiatic Society of Bengal* (Vol. LXVII, Part II, No. 1 ; 1898) settled this question, having noticed the change take place in a specimen which he kept, and which, when he got it, was mostly purple but by the first week of August was in non-breeding dress. We are inclined to agree with Finn as to the change and, from observations made during the current year (1924) at Pusa, can state definitely that males in full breeding (purple) plumage are not to be found during the Rains when the majority of males are in non-breeding plumage although an occasional one may be seen in a sort of half-and-half condition. The female is greenish-brown above, rather bright yellow below with the tail brown or blackish and the laterals narrowly tipped with white. The male especially is a most pugnacious little creature, not hesitating to attack



other birds much larger than himself, and sometimes even scrabbling and tapping at windows, apparently attempting to assault his own reflection in the glass. The breeding season in Bihar is from February to May, the earliest nest found with eggs in it being on 12th February and the latest on 30th May, and there are generally two or more broods in rapid succession, usually in the same nest. In Northern India, the laying season is later, in May and June, or as late as July or even August. In Southern India, eggs are to be found from January to June, but mostly in February to April. It breeds all over the Plains and to the summits of the Hills in Southern India and up to a height of about 5,000 feet in the Himalayas.

The nest is hung suspended from a twig or any convenient support and is composed of a very miscellaneous and heterogeneous mixture of materials beautifully woven together with the silkiest fibres and cobwebs, hair, fine grass, leaves, small pieces of dead wood, chips of bark, lichens, rags, scraps of paper, thorns, etc., all being made use of. The body of the nest is usually oval, with various scraps of material hanging below, while the apex of the oval is produced into a cone meeting the point of support. The nest is lined with silky-white seed-down, very neatly affixed to the interior. The entrance is a small oval or circular hole, about the centre of the nest and in some cases, but not always, a small projecting cornice is constructed just above the entrance; at Pusa practically all the nests seen have been provided with this little porch over the entrance, which does not seem to be oriented in any particular direction. The purpose of the porch is apparently to keep rain out of the nest. On the side opposite to the entrance the wall of the nest, which is ready some days before the eggs are laid, is pushed out a little so as to give room for the tail of the sitting bird. This pushing out of the back of the nest is one of the last portions of the work, and the female may be seen going in and out to try the fit, over and over again. When sitting, the head of the bird is seen just peeping out



of the entrance. The completed structure looks like anything but a nest and resembles a mass of dead leaves and rubbish caught in an old spider's web. The rows of sticks which support edible and sweet peas in our gardens form a very favourite nesting-site for this bird in North Bihar, as these sticks are in the ground during February and March, but the nest may be suspended almost anywhere, usually from the end of a branch of a bush or small tree, preferably a thorny one. A creeper, such as rose, *Quisqualis*, or *Stephanotis*, growing up the outer wall of a bungalow, is a very favourite site. It is usually placed between two and five feet from the ground but we have seen one as high up as forty feet. The nest shown in our Plate, as will be seen, was hung on to a piece of wire netting. Suspended in this way it is tolerably safe from enemies. Two is the normal number of eggs laid, but occasionally there may be three, the eggs measuring about 16 by 11·5 mm., being dull white marked with various shades of brown. Sometimes at least, only one of the young is reared up, and indeed there seems no room in this small nest for more than one fully-fledged nestling.

This bird, with other Sun-birds, is protected throughout the whole year in Burma, Bombay and Bengal, and presumably in Mysore also, as it is certainly a "bird of song."

Mr. C. M. Inglis has kept a specimen of the Purple-rumped Sun-bird in captivity for six months, feeding it on a mixture of sugar and *sattoo*.

Our Plate gives a good idea of the Purple Sun-bird and its nest. To the left is seen a male in breeding plumage with its tufts displayed; the female is seen feeding a young bird whose beak is seen projecting from the nest; and below her, to the right, is a male in non-breeding plumage. In Sind, Baluchistan and Afghanistan this Sun-bird is represented by a local race, the

Baluchistan Purple Sun-bird (*Cyrtostomus asiaticus brevirostris*),
and in Assam and Burma by the Burmese Purple Sun-bird
(*C. asiaticus intermedius*).



The Black-headed Oriole.



THE NORTHERN GOLDEN-BACKED WOODPECKER
(*Brachypternus aurantius aurantius*).



THE NORTHERN GOLDEN-BACKED WOODPECKER

(*Brachypternus aurantius aurantius*).

Numerous are the birds : woodpeckers sound
The barky trunks

Nebula to Man.

MANY of the roads on the Pusa Estate being well bordered by trees, it is a common experience to hear the quickly-repeated tapping sound of a woodpecker as it digs away with its beak at a branch in which it has perceived the presence of some wood-boring insect. If the noise is traced to its source, it will usually be found to originate from a medium-sized bird coloured in black and golden-yellow and with a bright crimson crest, which may be seen clinging to the bark with its powerful claws, supported by its short stiff tail, and hammering away with its short, stout, chisel-like bill. This bird is the Northern Golden-backed Woodpecker (*Brachypternus aurantius aurantius*), which is common throughout North India, Bombay, Central India, Bihar, Bengal and Orissa, ascending the hills to about three or four thousand feet, but not apparently known from Upper Assam or Burma or in the Duars. It is generally seen singly or in pairs and often one may be seen following the other from tree to tree. In Sind, Baluchistan, and parts of the Punjab it is replaced by the Sind Golden-backed Woodpecker (*B. aurantius dilutus*), in Southern India and part of Ceylon by the Southern Golden-backed Woodpecker (*B. aurantius puncticollis*) and in other parts of Ceylon by the Red-backed Woodpecker (*B. aurantius erithronotus*), which last, as its name implies, is distinguished by having its back crimson



whereas in the other races of *B. aurantius* the back is golden-yellow or orange. Where both sub-species occur together, it is probable that they hybridize occasionally. As regards other species of woodpeckers with which this one may be confused, the Common Golden-backed Three-toed Woodpecker (*Tiga javanensis*) is extremely like this bird and only markedly differs in wanting the hallux (inner hind toe). Tickell's Golden-backed Woodpecker (*Chrysocolaptes guttacristatus*) has also a very similar coloration but on account of its very much larger size is less likely to be mistaken for *B. aurantius* than is *Tiga javanensis*.

The Golden-backed Woodpecker is an extremely handsome bird, which Dewar briefly describes as having a bright crimson crests top of head black, sides of head white with a number of black line, and streaks, upper back golden-yellow, lower back and tail black, wings black and golden-yellow with some white spots. The female differs from the male in having the top of the head black with small white triangular spots; it is shown peeping around the tree in our Plate. Unfortunately, like the Indian Roller, its voice is not in harmony with its plumage, its call, which is often uttered on the wing, being a loud harsh scream. Like the Roller also, it is rather a noisy bird.

Like all woodpeckers, it is an extremely skilful climber, seldom or never perching crosswise on a branch but clinging, always with the tail downward, to the stems and branches of trees, which it usually ascends and descends diagonally. The flight is undulating. It is seldom seen on the ground but occasionally descends to feed on ants which seem to form an appreciable proportion of its normal food, this consisting almost entirely of insects, largely ants varied with numerous small beetles, caterpillars and bugs, to which buds and fruits may occasionally be added. The late C. W. Mason examined the stomachs of sixteen birds throughout the year at Pusa and found 3,921 insects of which the great majority were ants, and

it is notable that only one of these sixteen birds was found to contain longicorn beetle larvæ, although observation renders it certain that this bird does feed to some extent on wood-boring insects, which its long, worm-like, extensile tongue, armed with a many-barbed horny tip, is so admirably fitted to extract from their burrows after these have been laid open by the vigorous blows of the strong chisel-like beak. In Eastern Bengal it is stated to feed on the larvæ and pupæ of *Hoplocerambyx spinicornis*, a longicorn borer pest of *sal* (*Shorea robusta*), and in Madras it is stated to be very partial to toddy-palms, which may be due to the fact that these trees are infested by *Oryctes rhinoceros* and *Rhynchophorus ferrugineus*. At Pusa many of the dead *sissu* branches, at which one sees this woodpecker tapping away, are infested with a tree-living termite (*Coptotermes heimi*) and, although Mr. Mason's stomach-records do not support this, it is probable that this termite may provide a certain proportion of its food. As is the case with so many of our common Indian birds, it is wonderful how little we really know about its daily life. However, from the little we do know, we are justified in counting the Golden-backed Woodpecker, with all its kindred, amongst the farmer's friends.



Golden-backed Woodpecker.



Like other woodpeckers it nests in a hole in a tree, often in a mango tree in Northern India; in Bihar nests have been found in mango, litchi, *sissu* and *siris* trees. In the Southern part of its range it breeds from February to June but in the Northern part it breeds from March to July, courtship being of a rather rough and ready fashion, punctuated by harsh screams. The glossy white (delicate salmon-pink when fresh and unblown) eggs, usually three in number, measuring about 28 millimetres long by 20 broad, are laid in a hole excavated in a tree by the parents or more frequently in a natural cavity to which merely an entrance has been made by the birds. Dewar states that "woodpeckers seem to excavate a new nest every year," but we are not aware how far this is the case. Woodpeckers, however, are very apt to desert any nest-hole that has been interfered with at all. We have many times found them to do this even when no chipping was done to enlarge the hole, the only interference being by means of a thin twig. Sometimes also they only partly excavate a hole and then leave it for another site. The nest-hole is about two and a half to three and a half inches in diameter and usually runs in horizontally for about three to six inches and then turns downwards. When the downward shaft is bored by the bird it is rarely more than eight or nine inches deep with a chamber of some five or six inches in diameter, but when the bird cuts into a natural cavity in the tree the egg may be found two or three feet below the entrance. No regular nest is formed, the eggs being laid on a few chips of wood. The young, when first hatched, are naked but assume the sexual coloration with the first feathers. If caught young, they may be trained to a diet of "*sattoo*" with some soft fruit and occasionally some insects.

In Bengal, Burma, Madras, Bombay and Assam this bird is protected by law throughout the whole year.



THE NORTHERN GREEN BARBET
 (*Thereiceryx zeylanicus caniceps*)



THE NORTHERN GREEN BARBET

(*Thereiceryx zeylanicus caniceps*).

ALTHOUGH common enough where it occurs, the Green Barbet is not found in the less wooded areas of the country, such as the Punjab and Sind, Rajputana, and the more open parts of the Deccan and Carnatic. In areas where it does occur, however, it is sufficiently a common bird to merit inclusion in our list, especially as its call is probably more familiar than its personal appearance. It is a thick-set bird, about as large as a myna, with a thick, heavy bill, in colour of a bright leaf-green with a brownish head and a bare brown patch around the eye; but, as it always keeps near the tree-tops, it is less often seen than heard. Sometimes it may be seen on the wing, when its flight is strong but rather heavy and undulating. Its call is loud and monotonous, but by no means harsh or discordant, and is usually written *tur-r-r-r*, *kutur*, *kutur*, *kutur*, the call being indicated in the various vernacular names, such as *Kotur* (Hindustani), *Kuturga* (Maharatti), *Kotoruwa* (Sinhalese) and *Kutur* (Tamil). The call often sounds much like the word *Pakrao*, repeated several times.

Like most other birds, however, this Barbet is not confined to the use of a single note. In this connection Stuart-Baker, in his "Birds of North Cachar," writing about a very similar and closely allied bird, the Assam Lineated Barbet (*Thereiceryx lineatus hodgsoni*) makes the following interesting remarks: "I have heard this bird libelled as being a bird of one note; now any one who has listened to it *carefully* must admit that it has many. Whilst feeding, it has a large variety of sounds at its disposal. When



pleased, it utters a sort of hoarse 'chortle'; but, to make this sound, it seems to be necessary to be on the move, for it always utters it when hopping from one branch to another, or else it gives a little jerk into the air at the same time that it opens its mouth to give vent to its feelings. Displeasure, which seems to be caused chiefly by seeing other birds feeding with it, is expressed by a ridiculously feeble little sound like 'pench, pench,' the feebleness being made up for, to some extent, by the bird's ferocious attitude as he advances, with drooping wings and mouth wide open, towards the object of his displeasure. The most unusual note is one it makes use of only in the cold weather, at which time these birds sometimes collect in small flocks, and only in the mornings and evenings, seemingly for the purpose of collecting any scattered individuals. It consists of a loud clear whistle, a most wild and penetrating sound, but at the same time rather musical than otherwise. It is an abnormal sort of a sound for a barbet to give utterance to and had I not followed up and shot some of these birds whilst actually whistling thus, I should never have imagined what had made the sound." The Green Barbet may sometimes be heard calling at night, especially on moonlit nights.

Like other Indian Barbets, this bird lives chiefly on fruit. The late Mr. C. W. Mason examined the stomachs of fifteen birds at Pusa and found nothing but wild fig fruits in them. It is very fond of *Lantana* berries and helps to distribute the seeds of this noxious weed. This bird, however, has a curious habit of pulling off bits of bark from trees, especially from dead branches, as if searching for insects, and Blanford states that it is said occasionally, though rarely, to eat insects. Insects, however, evidently form a very small part of its diet and from an economic point of view this bird cannot be claimed as useful.

Blanford states that the characteristic "call is heard from January or February till June," and Dewar also says that this is



heard "during the latter part of the cold weather and the early part of the hot weather." Where the bird is common, however, the call may be heard throughout the year, although more persistently during the first half of the year. During the Rains, one of these birds may often be heard calling softly, as if talking to itself, and occasionally one or more birds give utterance to the usual loud call. As I write these lines (8th November) a Green Barbet is calling at intervals in a nearby tree.

Nesting takes place in a hole in a soft-wooded tree, usually in a dead branch or main stem, excavated by the bird itself, for which purpose its heavy bill seems well adapted.

The nest-hole is rather small, about five inches in diameter, and the passage leading into it is about six inches to two feet long, about two and a half inches in diameter, and very smoothly rounded off inside and bevelled off at the entrance, which is often situated on the underside of a branch and which in any case is always so placed that it does not face upwards, thus avoiding flooding of the nest-hole by rain. The nest is usually placed fairly high up in a tree, twenty feet or more above the ground, but may occasionally be found lower down. No regular nest is constructed, the eggs being placed on a few chips of wood in the nest-hole. When nesting, the birds are very shy about approaching or leaving the nest-hole whilst they are under observation.

In Bihar the nesting season is chiefly in March and April, but further North eggs may be found in May or even in June. The eggs, which are dull white, and slightly glossy, measure about 30 by 22 mm., and three or four eggs are usually laid.

The Green Barbet is now divided into three sub-species, the typical form (*T. zeylanicus zeylanicus*) being found in Ceylon and

South Travancore, the form found along the West coast of India from North Travancore to Bombay being *T. z. inornatus*, and that found in Northern India, from the extreme West to Western Bengal, being *T. z. caniceps*. It is this last sub-species which is represented in our Plate.

Another Barbet, which occurs along the lower ranges of the Himalayas and down to the South of Burma and which may often be seen in Calcutta, is the Blue-throated Barbet (*Cyanops asiatica*) which is a green bird, as large as a myna, with sky-blue patches on the cheeks and throat. Its call consists of a long series of thrilling notes, usually "Kürřawük, Kürřawük..." but sometimes "Kükärrük, Kükärrük....," preceded by a number of low, clucking notes. Like the Green Barbet, this bird nests in a hole which it excavates in a dead stump or branch of a tree.



The Blue-throated Barbet.



THE COPPERSMITH
 (*Xantholama haemacephala indica*).



THE COPPERSMITH

(*Xantholæma hæmacephala indica*).

Alone at his green forge
Toiled the loud coppersmith.

The Light of Asia.

ON the very hottest day of the year, when the brazen sky compels every animal to take advantage of any piece of grateful shade and the loquacity of even the crow is stilled, a soft, liquid, monotonous and sustained "tonk-tonk-tonk" gives evidence that the Coppersmith does not find the heat too enervating for song. Its note, which may be compared to the distant sound of a smith beating a sheet of metal, whence its popular name of Coppersmith, is sufficiently familiar during the hot weather in almost every garden throughout the Plains of India, but to most of those who hear its call it is indeed a case of "vox et præterea nihil," the owner of the voice being rarely observed unless especially looked for. Not that its appearance is anything to be ashamed of; on the contrary, it is a most gorgeously coloured little bird, dressed in green and crimson, the details of which will be seen on reference to our Plate.

The Coppersmith is usually referred to in books on birds as the Crimson-breasted Barbet and, as this name implies, belongs to the family of Barbets, which comprises upwards of twenty species of smallish, thick-set, green, fruit-eating birds, with a short strong bill, rounded wings and ten tail-feathers. We have already dealt with this group in connection with the Common Green Barbet (*Thereiceryx zeylanicus*) and it is sufficient to remark that the Coppersmith may be distinguished by its yellow throat and blackish



cheeks from the other species of *Xantholæma*, which are themselves separated from the other Barbets by the second primary being longer than the eighth, the lower tail-coverts green and the rictal bristles long.

In his fascinating book *Some Indian Birds and Acquaintances*, Cunningham has an excellent account of the Coppersmith, too long to quote in full, but from which we take the following extract:—
“Coppersmiths are odd little birds, and most fully characteristic of the group to which they belong in their gaudy colouring, harsh, dry plumage, wonderfully tough skin, and insistent vociferation. During periods of settled, sunny weather, the only thing that seems effectually to check their desire to call is a certain degree of cold, but this is so influential that during the course of the variable winter in Calcutta it may safely be assumed that the temperature in the shade stands at or over 70°F. on any day when their call is to be heard. As the thermometer rises above the prohibitive limit they begin to call more and more frequently until in the height of summer the monotonously metallic ringing of their notes goes on, almost constantly, from dawn to sunset. When preparing to call they usually take up a prominent place in the crown of a tree, often clinging to the side of an upright twig; and all the time that they cry they go on constantly turning their heads from side to side whilst their throats swell and their whole bodies thrill with the force of their vocal efforts. The movements of the head give rise to a strangely ventriloquial effect, so that the successive sounds might readily be mistaken for the answering notes of two birds instead of the continuous call of one. Towards the end of the hot weather, and during the early part of the rainy season, they cease to cry so incessantly, because the care of their young families takes up too much time to leave them much leisure for any other occupation.”
To some people the monotonous note, repeated indefinitely like the tick of a clock, is a source of exasperation until they are ready to exclaim with the apostle, “Alexander the Coppersmith hath



wrought me much evil," and in some parts of India which are not cursed with the ear-splitting shrieks of the Hawk-Cuckoo, the latter's name of "Brain-Fever Bird" is even at times misapplied to the Coppersmith; but to most people their song is soothing and it is at least constant and consistent and devoid of the exasperating quality of intermittence found in the true Brain-Fever Bird.

Like other Barbets, the Coppersmith is a frugivorous bird and seems to confine itself entirely to wild fig fruits, not being known to attack cultivated fruits at all.* The late C. W. Mason examined the stomach-contents of fifteen birds at Pusa and Mr. D'Abreu those of two more at Nagpur, and in all cases nothing but *Ficus* fruits was found to have been eaten. From an agricultural point of view, therefore, this bird is of neutral value.

The nesting season is in the early part of the hot weather, from March to May in Northern India, earlier in Central and Southern India. The nest, if such a term may be used, is placed at the end of a long gallery, which may be from one to five feet in length, either excavated by the parent birds in sound though soft wood of a branch, or more frequently in a branch which is already decayed internally and into which the birds cut, through the harder external shell of the branch, a perfectly circular hole with the edges neatly bevelled off inside and outside, this hole being about two inches in diameter and always placed on the underside of the bough. If a ready-made hollow is available it is preferred and appropriated, but if the birds have to excavate a new gallery on their own account, they usually choose a place on the under surface of a slanting dead bough, especially at a point where a side branch has been broken off

* Mr. C. M. Inglis, however, writes that prior to painting the Plate he carefully studied the bird so as to try to obtain a natural position, and one morning, whilst watching several of them on a guava tree in fruit, he saw them eating voraciously, digging out huge holes in the ripe fruit. There were about half-a-dozen birds hard at it at the same time and they played havoc with the tree. Mr. Inglis has also seen these birds sometimes hawking for termites. [T. B. F.]

and the wood softened by the invasion of fungal mycelium. Whilst working, the bird clings to the bark like a woodpecker, with the end of its tail pressed closely against the surface, so as to serve as an additional support whilst it hammers and picks away with its bill at the soft wood, the chips of wood being sometimes hammered directly off but oftener merely loosened by a series of blows and then picked up and thrown away. Both of the parent birds take part in the work of excavation and they can sometimes be heard excavating their holes long after sundown. The eggs are placed at the bottom of the cavity into which they have thus bored, and which they smoothen a good deal interiorly, often a couple of feet below the entrance, and laid merely on the chips produced in the course of the work. Three or (more frequently) four eggs are laid, the egg being on an average 25 mm. long by 17.5 mm. broad, delicate pink when fresh, pure white later, almost cylindrical, tapering somewhat towards one end, but with the ends themselves broad and obtuse. The young birds are much like the adult, but duller in colour and without the crimson and black markings on the head. Occasionally specimens occur in which the green plumage of the adult bird is replaced by yellowish.

The Coppersmith, we regret to say, seems rather an ill-natured little bird and is therefore not adapted to be a desirable inmate of a mixed aviary. He is not social with other birds rather than actively aggressive, but requires plenty of space and must be fed on his natural fruit diet, as *sattu* (pea-meal paste), although eagerly devoured by this bird, is rapidly poisonous to it.



Brain-Fever Bird.



THE NORTHERN INDIAN ROLLER

(*Coracias benghalensis benghalensis*).

Decked with diverse plumes, like painted jayes.

The Faërie Queen.

Entrancing 'tis to watch their turquoise flight

With wings extended

Chenar Leaves.

Undreamed-of wings he lifted.

THE Indian Roller or Blue Jay (*Coracias benghalensis* formerly known as *C. indica*) is one of the birds of the Plains which must have brought itself to the notice of all if merely on account of its vivid colouring, which has been aptly described as a study in Oxford and Cambridge blue. Although commonly called a jay, this bird is more nearly allied to the bee-eaters and kingfishers, its relationship to the latter group being evidenced, as recorded by Gordon Dalglish, by the fact that it has been seen to plunge into the water like a kingfisher, this most unusual occurrence indicating an affinity to the kingfisher in habits, especially to the white-breasted species which is practically an insect-feeder. As a matter of fact, *C. benghalensis* belongs to the group of Rollers, so called because of the extraordinary aerial gymnastics indulged in by these birds, as may be seen especially during the early part of the hot weather, when courtship takes place.

The Indian Roller occurs throughout the Plains of India and Ceylon, neither ascending the Hills nor occurring in areas of desert or thick jungle. The form found throughout Northern India to Eastern Bengal is the Northern Indian Roller (*C. benghalensis benghalensis*), which is replaced in Southern India and Ceylon by the

Southern Indian Roller (*C. benghalensis indica*). Calcutta seems to be about the eastern limit of occurrence of the Northern sub-species, which East of that is replaced by the Burmese sub-species,



Indian Roller

Coracias benghalensis affinis, which is slightly larger and much darker, with a lighter tail which lacks the purple band at the tip which sets off so well the tail of the Indian Roller. In the Duars both species have been got and where they meet they hybridize



freely. In Northern India another species, the Kashmir Roller, *Coracias garrula semenowi*, a local race of *C. garrula*, which is found in Africa and Europe, also occurs and may be distinguished by its lower parts being pale blue throughout, whereas in the Indian and Burmese Rollers, the lower parts are only blue in part. There is also a forest species, *Eurystomus orientalis*, the Broad-billed Roller, with a red bill and legs; this is a rather silent bird and exceedingly wary and frequently nests in non-accessible holes in large *Simul* (*Bombax malabaricum*) trees.

Where it does occur, the Indian Roller is usually found commonly and is not a bird which can be overlooked when on the wing, as its brilliant blue colours are then displayed to advantage. When at rest upon a branch or telegraph wire, however, this bird is by no means conspicuous, as the wings are then closed and the colours are not very evident. From the sudden contrast between the inconspicuous hues of this bird when at rest and the vivid display of colour evidenced when it takes to wing, it is sometimes known as the "Surprise Bird." Its usual habit is to sit on any convenient perch and watch patiently until some desirable prey comes into sight, when it flies down, secures its quarry, which is always swallowed whole, and returns to its perch. Ordinarily it appears to be a decidedly sluggish bird, sitting for hours on the same perch, occasionally jerking its tail and emitting a sharp harsh "Tjock." The extremely inconspicuous appearance of this bird when at rest provides concealment at once from enemies and from prey and the Roller seems to be well aware of this fact, as, if it has no reason to suppose itself to have been noticed, it will often permit very close approach, remaining quite motionless on its perch. To a potential enemy, which discovers it at close quarters, the sudden exhibition of its brilliant colouring must have a startling effect.

At the breeding season, which occurs in Bihar from March to the end of June, and as early as January in Ceylon, these birds, however,



become more active and vociferous, their harsh, staccato gutturals being evidently pleasing to the opposite sex. It is at this time also that they indulge in the weird evolutions which have earned these birds the title of Rollers. As both sexes wear the same livery it is difficult to tell which is which, but two birds may often be seen sitting side by side on some exposed perch during the breeding season and uttering a sort of chuckling sound. One of them, presumably the cock bird, then flies off and up into the air to a considerable height whence he descends in a regular "nose-glide," displaying his vivid colours and uttering short harsh screams all the while, ultimately returning to perch beside his mate.

The nest is placed in a hole in a tree or building and is generally lined with a varying amount of vegetable fibre, grass, a few feathers, or some old rags, but the lining is often omitted altogether. Four, or sometimes five, glossy-white eggs are laid. The young are quite naked at first, but, later on, when the feathers have grown, are marked with vivid blue like their parents. Mr. Aitken considered the parent birds very wary and the nests difficult to discover, but in Bihar at any rate this is not the case, it being one of the easiest nests to find, the birds continually chasing away any intruders that come near the nesting site.

Besides being, as the poet puts it, "a lovely bird with azure wings," and thus an ornament to any landscape, the Indian Roller is an extremely useful bird, as its food consists almost entirely of large insects, such as grasshoppers, crickets and beetles, varied by an occasional small mouse, frog or snake. One kept by Mr. Finn even digested a toad, which must have been a very "tasty" morsel. The late C. W. Mason examined the stomachs of eighteen birds between January and October at Pusa and found that of 412 insects taken by these birds, only 4 were beneficial, 111 were injurious and 297 neutral. Of the injurious insects taken 52 were grasshoppers, 18 crickets and 23 caterpillars, mostly cutworms, so



that the good done by the destruction of these injurious insects far more than counterbalances the fact that a very few beneficial insects were taken.

Towards the end of the Rains, when the big brown crickets emerge at dusk from their holes in the soil, the Roller is well aware of their habits and may be seen, until night has almost fallen, perched on a convenient bough and watching eagerly for the chance of a tasty supper, a happy chance which is accomplished should any unwary cricket wander far enough to enable it to be captured before it can regain the safe security of its burrow. There is in my garden a large *Dalbergia* tree with a dead branch which affords an excellent clear view of the ground beneath, and every evening at sunset a Roller takes up its post on this dead bough and watches until far into the dusk for the crickets which are foolish enough to leave their burrows in the lawn and flower-beds beneath this tree. The Little Owl, likewise fond of crickets for supper, also appears at the same time but, having the advantage of nocturnal vision, is able to harry the crickets throughout the night as they forage for food. Like Owls also, the Roller throws up the undigested portion of his food in the form of pellets.

It is therefore unfortunate that the Roller's brilliant plumage should frequently lead to its being shot by Europeans, especially in the neighbourhood of military cantonments, for the sake of its wings. According to the late C. W. Mason, "being one of our common species of birds, and the gaudy colour very striking to any one new to the Country, numbers of these birds are shot by Europeans in order to send one or two wings home, and they are sent home not declared, or falsely declared. From what I have seen, I do not imagine that more than one out of six pairs of wings ever sees the destination for which they were originally obtained. Some specimens are not good enough, while others are put away, forgotten, and eventually thrown away." It should therefore be


noted that the Indian Roller is protected, under the Wild Birds and Animals Protection Act, throughout the whole year in Bihar and Orissa and in Delhi. In view of its beneficial activities to the Indian *raiya*, it is to be hoped that the numbers of this bird will not be allowed to diminish throughout the area of its occurrence.

This bird is sacred to Vishnu, who once assumed its form, and is caught to be liberated at the Hindu Dasahra festival in Western India and at the Durga Puja in Bengal. Finn states that "once at least a throne has been gained by the holy fowl—no doubt a trained specimen—alighting on the successful candidate's head."

Falconers fly the Red-headed Merlin (*Æsalon chicquera*) at this bird. Jerdon says "it is often baulked by the extraordinary evolutions of the Roller who now darts off obliquely, then tumbles down perpendicularly, screaming all the time, and endeavouring to gain the shelter of the nearest tree or grove."



The King Crow.



THE COMMON INDIAN BEE-EATER
(*Merops orientalis orientalis*).



THE COMMON INDIAN BEE-EATER

(*Merops orientalis orientalis*).

Nor hawked the merops, though the butterflies—
Crimson and blue and amber—fitted thick
Around his perch.


The Light of Asia.

THE Common Indian Bee-eater is a small slenderly-built bird, of a bright green colour, with a rather long curved black beak and with the two middle tail-feathers projecting like bristles a couple of inches beyond the other tail-feathers; the eye is bright red, the throat green or blue (whereby it is readily distinguished from the Blue-tailed Bee-eater with its chestnut throat), a black band under the eye and a black collar above the breast, some black in the tail, and the lower surface of the wings bronzy-red. It is easily recognizable and not likely to be confused with any other bird found commonly in the Plains. There are three sub-species, *Merops orientalis orientalis*, whose distribution is given as India, Bengal to Ceylon; *M. orientalis birmanus*, the Burmese Green Bee-eater, found in Assam and Burma; and *M. orientalis biludschicus*, the Sind Green Bee-eater, which ranges from South-East Persia to Sind and Baluchistan. These sub-species vary in the amount of red-gold about the head, this being wanting in the Sind birds and more pronounced in the Burmese race.

Cunningham, whom we have quoted so often, says of this bird:—"Common Bee-eaters are singularly alluring both in appearance and in the character of their notes. When seen from behind they look brilliantly green with golden gleams about their heads;

their wings have a ruddy bronze tint, and there is a beautiful patch of blue on the throat—a scheme of colour which, along with their brilliantly sparkling eyes, cheerful cries, and confiding familiarity, is well adapted to command general admiration.”

This Bee-eater occurs throughout India and is usually abundant when it occurs, but it seems to be a local migrant, moving to different localities in the hot and cold seasons, so that in some



Common Bee-eater.

places it may be found only at one season and in other places, where it is seen all the year round, it is probable that the individuals seen (with the exception of a few remaining to breed) do not remain in that place all the year. There seems to be a regular movement of the species southward about September and northward again about March in every year. Thus, this bird is found in Madras in the cold weather but, with the exception of a few which remain



throughout the year, most individuals move north about March. In Calcutta the annual immigration takes place about the second week of October and the exodus about March, as soon as the hot weather has set in. In North Bihar this Bee-eater appears in numbers about the beginning of October and again in February and March, these temporary accessions apparently indicating the passing through of a migrant stream southwards and northwards respectively. In the Punjab and North-West Province, this bird arrives about March and leaves again about September. It seems probable that this migration is due primarily to the necessity for moving away from localities in which the necessary insect-food is becoming scarce, but we really know very little about the underlying causes of migration in birds. Cunningham says "Their departure can hardly be determined by dietetic causes, as other kinds of insectivorous birds continue to find an abundance of insect-food all through the summer. It is apparently due to their nesting habits, for, nesting as they do in burrows in the soil of fields and banks, in a region like the lower Gangetic delta, they must naturally meet with great difficulties in finding sites secure from repeated inundation during the torrential falls of rain that frequently take place during the summer months. Hence they move off to somewhat higher and dryer regions, and remain there until, with the colder and dryer days of autumn, insect-food becomes inconveniently scarce, and they are once again driven back to milder and damper places." But the last sentence of this quotation seems to be in direct contradiction to the first, nor does Cunningham's theory fit the case of many parts of Madras, where the rainfall is practically all during the North-East monsoon, in October and November. The fact seems to be that this Bee-eater prefers a dry, sunny climate, because such conditions are necessary for the capture of its prey which consists almost wholly of flying insects taken on the wing. Thus, it is not found in dense or damp forests and in Ceylon it only occurs in the drier parts of the low country.



The Common Bee-eater is a sprightly little bird with a loud but rather pleasant note. It is fond of resting on projecting boughs, railings, telegraph or other wires, or in any similar position affording a good view, whence a sharp look-out is kept for any insect seen moving on the wing, whereupon the bird projects itself into the air with a few quick strokes of the wings, then glides onwards with widely-spread wings and tail to secure its prey, and then usually returns to its perch to devour it. The snap of the beak with which the prey is caught is often audible from a distance of several yards. If the prey is small it may be swallowed at once, with a quick upward jerk of the beak, but if large or a stinging insect it is generally broken up a good deal by striking the lower mandible against the bird's perch; as these birds are usually very tame, it is often possible to watch this process at quite close quarters.

The food is composed exclusively of insects, in the broad sense of the word, and the largest proportion of it is composed of flying insects taken on the wing. Of these, bees and other Hymenoptera form the vast majority and this Bee-eater is a very serious pest in most localities in the Plains where apiculture is practised. Our apicultural experiments at Pusa have been largely brought to a standstill, almost wholly on account of the activities of these Bee-eaters, practically all the newly-raised queen bees being snapped up by them whilst on their marriage-flight. Dewar, in his *Birds of the Plains*, says, "I doubt whether the little *Merops viridis* tackles an insect so large as a bee." Our records show most emphatically that this Bee-eater not only tackles bees but is a perfect pest of the apiary, destroying very large numbers of honey-bees of all sorts. We have found at Pusa that it eats *Apis florea* (the small honey-bee), *Apis indica* (the Indian honey-bee) and *Apis mellifera* (the European honey-bee), whilst Mr. D'Abreu found at Nagpur that this bird had even eaten *Apis dorsata* (the large rock honey-bee). Out of forty-three birds' stomachs examined at Pusa by the late C. W. Mason, of 376 insects taken no less than 300 (mostly



bees) were classed as beneficial, 53 as injurious (mostly weevils and none of them real pests), and 23 as neutral. Some birds examined at Nagpur by Mr. D'Abreu contained honey-bees, dragonflies, blue-bottles, coprid beetles and one grasshopper. Mr. Inglis watched a bird at close range for twelve minutes during which it made twelve sallies after insects and it appeared to him that nine out of the twelve insects taken were bees. Besides bees, dragonflies are taken very largely; the species is not given as a rule but at Pusa I have seen this bird catch and eat *Potamarcha obscura*. So far as its feeding-habits go, therefore, this bird does not seem to deserve the protection throughout the whole year which it enjoys at present under the Wild Animals' Protection Act in Bombay, Bengal, Assam and Burma. It seems a pity that such a lovely little bird should be so noxious in its habits, "but pity 'tis, 'tis true."

The Common Bee-eater breeds from the middle of March to the beginning of June, excavating a long gallery, generally four or five feet long, in a dry sandy bank. At Peshawar I have found it nesting in May in holes dug in sandy soil near the top of a river bank. As a rule, this bird does not seem to breed much south of the Punjab but a few individuals remain in most localities, even as far south as Madras, and breed there. The excavation of the gallery for the nest is done, in alternative short spells, by both the parent birds, and when the gallery, which is about two inches in diameter, has been carried far enough within the bank, its end is widened out into a rounded chamber about twice the diameter of the gallery and in this chamber, without any further attempt at a nest, the eggs are laid to the number of three to five, usually three or four, occasionally more. Mr. Inglis has taken up to seven eggs and has also found a single young bird in a nest. The egg is a spherical oval, white and glossy, about 19 mm. by 17 mm. A nest-hole is shown in our Plate, which gives a good idea of this bird.

Blanford gives the following vernacular names of the Common Bee-eater, *Patringa* and *Harrial* in Hindi, *Banspati* in Bengali, *Tairlingi*, *Veda Raghu* in Mahratti, *Chinna passeriki* in Telugu, *Kattalan Kuruvi* in Tamil, *Monagyi* in Arrakan, and *Hnet-pasin-to* in Burmese ; but, as he remarks, several of these terms are applied indiscriminately to other species of Bee-eaters also.



A female Honey-sucker.



THE BLUE-TAILED BEE-EATER
(*Merops superciliosus javanicus*).



THE BLUE-TAILED BEE-EATER

(*Merops superciliosus javanicus*).

In the mango-sprays bee-eaters hawked
Chasing the purple butterflies.

The Light of Asia.

To my fair garden, clad in verdure green,
They come at times, perchance to look for Spring,
With glowing tints of blue and emerald green
On downy breast and softly-spreading wing,
And o'er the glories of my flower-bed
Flash by like sunbeams dancing overhead.

MANY of the previous papers have dealt with useful birds—useful, that is, so far as agriculture is concerned—but the same adjective cannot be applied to the Blue-tailed Bee-eater, as it is on the whole an injurious species by preying on beneficial insects.

In appearance and habits it is much like its commoner and smaller, but equally noxious, relative, the Common Indian Bee-eater (*Merops orientalis orientalis*), from which it is distinguishable by its larger size and by its chestnut throat (green or bluish-green in *Merops orientalis orientalis*). It is shortly described by Dewar (*Indian Birds*, p. 161) as “general hue green, shot with bronze; the tail is bluish. There is a broad, black streak running through the eye. The chin is dirty cream colour. The throat is chestnut-red. The eye is bright red.”



The Blue-tailed Bee-eater occurs commonly throughout the Plains of India but is partially migratory, visiting Northern India in the summer and Southern India in the winter. In the Central Provinces it is stated by Mr. D'Abreu to be resident throughout the year. In Bihar they are seen from March to October and in the Duars have only been noticed during June and July. Mr. Stuart-Baker only once came across it in North Cachar. Usually it is seen in small numbers or singly in one place but occasionally it congregates in large numbers. As a rule it is commonest in well-wooded districts. It is fond of perching on a post or telegraph wire or other suitable situation whence it swoops down on its prey which is usually captured on the wing, the bird thereafter returning to its perch. The flight is swift and graceful and the note a mellow whistle continually uttered while on the wing. During evenings, especially after rain, numbers may be seen hawking about on the wing for a long period without settling.

In Calcutta the Blue-tailed Bee-eater appears during the Rains, usually in large flocks. It is often to be seen in large numbers in places abounding with open sheets of water, but it is also met with not infrequently in gardens where its elegant flight makes it a decided ornament. It is, however, a shyer bird than the Common Bee-eater and its habits are not so easily observed at close quarters. It has been noticed to hawk its insect prey just over the surface of the water and sometimes even to pick an insect off the surface of the water.

This Bee-eater is common at Pusa and the late C. W. Mason examined the stomachs of thirteen birds between April and October. He states that, of 83 insects taken, 70 were beneficial, 3 injurious and 10 neutral. The beneficial insects taken included dragonflies, honey and other bees, and wasps, and in the neighbourhood of an apiary these birds may be a decided pest by snapping up the bees. Mr. Mason noted that, on dull days when dragonflies are inclined



to sit on grass-stems, this Bee-eater feeds almost entirely on these insects and that the dragonfly, *Crocothemis servilia*, is one species which is taken in large quantities at such times.

Breeding occurs, usually in large colonies, some time between March and June, in a hole which may be four to seven feet long with a diameter of two to two and-a-half inches and which is excavated in a bank, usually a river bank though not always so, the egg chamber being about half a foot in diameter and, what is unusual with Bee-eaters, it is sometimes lined with grass or feathers. Four or five and rarely seven glossy, white, almost globular, eggs are laid. Our Plate shows the entrance of a nest.

Besides the present species there are six others found in India and Burma, viz., the Common Indian Bee-eater (*Merops orientalis orientalis*), a smallish green bird with a golden hue on the crown of the head and the usual black band under the eye. This is the commonest of all the Bee-eaters and as a rule resident wherever it occurs. We have already dealt with it. The Blue-cheeked Bee-eater (*Merops persicus*) is very like the Blue-tailed bird but has the upper surface of the tail green instead of blue. This is a migrant from Africa and West and Central Asia to the North-West of India, breeding there. The European Bee-eater (*Merops apiaster*), a smaller species with a yellow instead of a chestnut throat, is also a migrant from Africa, visiting Kashmir and the Punjab during May and June. The Chestnut-headed Bee-eater (*Melittophagus erythrocephalus*) has a short tail and is chestnut above: this species is mostly confined to forests and is found in Ceylon, Southern India, the lower ranges of the Himalayas, and in Burma. The Blue-bearded Bee-eater (*Nyctiornis athertoni*) and the Red-bearded Bee-eater (*N. amictus*) are stout birds for Bee-eaters, and have, as can be seen by the names, blue and red plumes down the breast; they are green above; these birds are not so graceful in their movements as the other Bee-eaters. Blanford says the nidification of the latter



bird is unknown but since then the eggs have been taken in Perak in February; they appear to be smaller than those of the Blue-bearded species.



Common Bengal Bulbul.



THE INDIAN HOOPOE
(*Upupa epops orientalis*)



THE INDIAN HOOPOE

(*Upupa epops orientalis*).

Hoopoes—whose frequent notes sound as their name—
A merciless toll on worms and insects claim.


Nebula to Man.

The hoopoe too, in fascinating crest
And wings all striped in pattern alternate—
Of white and black, will take among the rest
Of Kashmir's birds a place of honour great.
Hark ! how his name hoopoe will reproduce
His hollow notes in quaint similitude.

Chenar Leaves.

EXCEPT in Sind and the Western Punjab, the Indian Hoopoe is one of the most familiar of the birds which live in the Plains of India, occurring commonly, and usually abundantly as a resident in all suitable localities throughout India, Burma and Ceylon. It is a bird about the size of a myna, fawn-coloured, the wings and tail white with very broad black bars, the legs short, the bill very long and slightly curved, and the back of the head with a conspicuous crest, which is normally kept folded so that it looks like a backward prolongation of the bill, but which is erected like an expanded fan when its owner is excited or disturbed or flies and when it first settles after flight. Specimens from South India and Ceylon run smaller than those from Northern India and have been separated under the name *Upupa ceylonensis*. Burmese specimens run larger in size, and have sometimes been separated as a distinct species, *U. longirostris*, a name derived from the greater length of the bill in Burmese examples. Besides the Indian species, the European Hoopoe (*U. epops epops*), distinguished by the presence of white between the buff ground-colour and the black tips of the feathers of the crest on

the head, is found in the Himalayas during the summer and in the winter visits the northern half of India, extending as far South as the Deccan, Chota Nagpur and Sylhet. Some Indian specimens, especially those from the North, very often show a tinge of white on



Head of European Hoopoe (after Blanford).

the crest, and these are regarded as hybrids between the Indian and European species. In North Bihar many specimens intermediate between the Indian and European forms are to be seen. In most parts of India, however, the buff-coloured bird which is seen probing the lawn with its long bill may safely be put down as the Indian Hoopoe. A true albino with pink eyes has been recorded.

The Hoopoe is found chiefly in open country and is essentially a ground bird, only occasionally perching on trees. Its flight is slow and undulating. Like other birds which have developed a special type of bill, this organ is intimately adapted to its owner's method of obtaining food, which, in the case of the Hoopoe, consists of insects and occasional worms, obtained mostly on or from under the ground, rarely on trees or in the air. The major portion of its food is obtained either by probing grassland for caterpillars living at the roots of the grasses, or by turning over leaves and rubbish for insects. It rarely



picks them off plants or trees and still more rarely catches them on the wing, although winged termites are occasionally taken in this way. The favourite haunts of this bird are avenues, especially if grass is growing in these grasslands and lawns. It prefers slightly damp, but not wet, localities to dry ones, as its insect food is then more easily procurable. It is an interesting sight to see these birds regularly quartering a lawn, stopping every now and then to dig and probe the soil with their long beaks, the result of investigation generally yielding some insect, which is extracted and swallowed. If, however, the parent bird is collecting material to feed its young, it is often robbed by a King-crow, as we observed in our previous article on that bird.

As Mason remarks, the young birds are fed almost entirely on caterpillars (probably all cutworms), grubs of Melolonthids, and crickets, and the amount of food fed during the day to a nest of half-grown young is extraordinary. Mr. Mason watched a nest one day from 6 a.m. and in the first hour fifty-eight visits were made to the nest by the old birds, and during these visits forty-five insects were almost certainly cutworms, ten were other caterpillars and grubs (some almost certainly *Anomala* grubs) and three were crickets (one of these may have been a large beetle); during the next half-hour twenty-seven caterpillars and grubs were brought and fed to the young birds. All this food was obtained from grass lawns or under *Sissu* and mango trees. Only one insect was brought at each visit and all these insects were large ones. The same nest was watched again by Mr. Mason a week later, but the birds now seemed more wary and consequently but few of the insects fed could be identified but they appeared to be the same kinds as observed before. On the latter occasion 286 visits were paid to the nest by the parent birds between 6 a.m. and noon and about two visits per hour were made when apparently no insects were brought, so that about 274 insects were brought in and fed in six hours to the young birds, who



numbered four or five at most. The food brought to the young consists practically wholly of caterpillars, beetle grubs and crickets. When bringing in food to the nest, the old bird as a rule perches near the nest to look around for danger and almost always utters a harsh grating sound on approach to the nest and again on entering and leaving the nest.

The nesting season is in the early spring or beginning of the hot weather, and nesting takes place in any convenient hole, never at any great elevation from the ground, in trees, walls or banks. A nest has even been found on the floor of a house amongst some *bhusa* (chaff). The nest itself is a mere apology, a little hair, a few feathers, leaves or grass-stems being carelessly strewn over the floor of the hole or hollow and, when eggs are laid in a tree-hole, there is often no nest at all. These birds never remove the droppings from their nest and the stench of these is most overpowering. Four to six eggs are laid as a rule, but as many as nine are reported to have been met with occasionally. So many young are not always reared and one of these birds has been seen dropping two out of four of its young from the nest, presumably to reduce the number of mouths to feed. The egg is a very lengthened oval, pointed at one end and sometimes tending to be pointed at the other end also, not glossy, uniformly pale greyish-blue or olive-green or olive-brown or any intermediate shade, the average size being 24 by 16.5 mm. The female bird alone incubates the eggs and, especially when the eggs are near hatching, scarcely ever leaves them alone for a moment, being assiduously tended by the male bird, who brings her food continually. Mr. Inglis has seen a male bird, before the breeding season had commenced, run up to his mate and present her with an ant-lion grub. When the female bird is sitting closely in this way, she hisses like a snake if disturbed. They are very loath to leave the localities where they breed. On one occasion, to inspect a nest, the hen bird had to be pulled out of the hole and in doing so some



of her tail feathers came out, but even this rather severe handling did not make her quit the place.

The young nestlings, as noted above, are fed by the parents on caterpillars and crickets and, as soon as they are able to leave the nest, they may be seen trotting after their anxious parents, making inefficient attempts at digging on their own account, but always ready to run up and have supplies thrust far down their throats by the long, curved beaks of their guardians.

It will readily be understood that a bird which feeds on insects, as does the Hoopoe, is a very useful one to the farmer. From actual examination of the stomach-contents of twenty-four birds at Pusa, the late C. W. Mason found that these had swallowed 278 insects of which the majority belonged to injurious species. A large proportion of its food also consists of cutworms and other insects living below the actual surface of the ground, so that they are fairly immune to most other enemies, and from this point of view, as a destroyer of cutworms and cockchafer grubs, the Hoopoe is most decidedly amongst the farmer's best friends and deserves every encouragement and protection. It is protected throughout the whole year, under the Wild Birds Protection Act, in Bombay, Delhi, the United Provinces, Bihar, Bengal, Assam and Burma, but in Madras in the Shevaroy Hills only. In Mysore it is not specifically protected but is presumably included in the schedule which includes all birds of bright plumage.

Both the Indian and European Hoopoes are known in Hindustani as *Hud-hud* and in Mahratti as *Sutar*. The name *Hud-hud*, as also the English name Hoopoe, are both derived from the call of these birds, a gentle " ŪK, ŪK, ŪK, ŪK, ŪK," usually uttered when sitting on an exposed branch of a tree, the head being depressed until the tip of the beak almost touches the breast, the crest at the same time being laid flat down. This cry should not be confounded with the much louder deeper call of the Crow-pheasant.



Hoopoes have been kept in confinement and of course require an aviary and not a cage: the aviary should have turf laid down, as otherwise the bird's bill gets damaged whilst probing about for insects. Butler says "the best food for it consists largely of soaked ants' cocoons, supplemented by mealworms, spiders, insects of all kinds, and earthworms." They are said to get very tame in captivity.

The specific status of the Indian Hoopoe seems to be rather doubtful. In the third *Fauna* volume on Indian Birds, by Blanford, it is called *Upupa indica*, under which the named forms *ceylonensis*, *nigripennis* and *longirostris* are sunk, although the name *ceylonensis* has priority over *indica*, and the European Hoopoe is separated as a distinct species under the name *U. epops*. It seems doubtful how far the Indian and European birds are really distinct and it is probably better to include them all under the name *Upupa epops*, with the names *U. epops epops* for the European race, *U. epops orientalis* for the North Indian race, *U. epops ceylonensis* for the South Indian and Ceylonese form, *U. epops longirostris* for the form found in Burma, and *U. epops saturata* for the Tibetan form which is a winter visitor to Assam and Burma.

The Hoopoe is looked on as a favourite because, so the story goes, he was King Solomon's messenger, and he is known as the King of Birds from the legend of his crest and crown, which is related by the Hon. Robert Curzon as follows:—

In the days of King Solomon, the son of David, who, by the virtue of his cabalistic seal, reigned supreme over genii as well as men, and who could speak the languages of animals of all kinds, all created beings were subservient to his will. Now, when the king wanted to travel, he made use, for his conveyance, of a carpet of a square form. This carpet had the property of extending itself to a sufficient size to carry a whole army, with the tents and baggage; but at other times it could be reduced so as to be only large enough



for the support of the royal throne, and of those ministers whose duty it was to attend upon the person of the sovereign. Four genii of the air then took the four corners of the carpet, and carried it with its contents wherever King Solomon desired. Once the king was on a journey, in the air, carried upon his throne of ivory over the various nations of the earth. The rays of the sun poured down upon his head, and he had nothing to protect him from its heat. The fiery beams were beginning to scorch his neck and shoulders, when he saw a flock of vultures flying past. "O vultures!" cried King Solomon, "come and fly between me and the sun, and make a shadow with your wings to protect me, for its rays are scorching my neck and face." But the vultures answered, and said, "We are flying to the north, and your face is turned towards the south. We desire to continue on our way; and be it known unto thee, O king! that we will not turn back in our flight, neither will we fly above your throne to protect you from the sun, although its rays may be scorching your neck and face." Then King Solomon lifted up his voice, and said, "Cursed be ye, O vultures! and because you will not obey the commands of your lord, who rules over the whole world, the feathers of your neck shall fall off; and the heat of the sun, and the cold of the winter, and the keenness of the wind, and the beating of the rain, shall fall upon your rebellious necks, which shall not be protected with feathers, like the neck of other birds. And whereas you have hitherto fared delicately, henceforward ye shall eat carrion and feed upon offal; and your race shall be impure till the end of the world." And it was done unto the vultures as King Solomon had said.

Now it fell out that there was a flock of hoopoes flying past; and the king cried out to them, and said, "O hoopoes! come and fly between me and the sun, that I may be protected from its rays by the shadow of your wings." Whereupon the king of the hoopoes answered, and said, "O king! we are but little fowls, and we are not able to afford much shade; but we will gather our nation



together, and by our numbers we will make up for our small size." So the hoopoes gathered together, and, flying in a cloud over the throne of the king, they sheltered him from the rays of the sun. When the journey was over, and King Solomon sat upon his golden throne, in his palace of ivory, whereof the doors were emerald, and the windows of diamonds, larger even than the diamond of Jemshea, he commanded that the king of hoopoes should stand before his feet.

"Now," said King Solomon, "for the service that thou and thy race have rendered, and the obedience thou hast shown to the king, thy lord and master, what shall be done unto thee, O hoopoe ? and what shall be given to the hoopoes of thy race, for a memorial and a reward ?"

Now the king of the hoopoes was confused with the great honour of standing before the feet of the king ; and making his obeisance and laying his right claw upon his heart, he said, "O king, live forever ! Let a day be given to thy servant, to consider with his queen and his counsellors what it shall be that the king shall give unto us for a reward." And King Solomon said, "Be it so."

And it was so.

But the king of the hoopoes flew away ; and he went to his queen, who was a dainty hen, and he told her what had happened, and desired her advice as to what they should ask of the king for a reward ; and he called together his council, and they sat upon a tree, and they each of them desired a different thing. Some wished for a long tail ; some wished for blue and green feathers ; some wished to be as large as ostriches ; some wished for one thing, and some for another, and they debated till the going down of the sun, but they could not agree together. Then the queen took the king of the hoopoes apart and said to him, "My dear lord and husband,



listen to my words ; and as we have preserved the head of King Solomon, let us ask for crowns of gold on our heads, that we may be superior to all other birds."

And the words of the queen and the princesses, her daughters, prevailed ; and the king of the hoopoes presented himself before the throne of Solomon, and desired of him that all hoopoes should wear golden crowns upon their heads. Then Solomon said, " Hast thou considered well what it is that thou desirest ? " And the hoopoe said, " I have considered well, and we desire to have golden crowns upon our heads." So Solomon replied, " Crowns of gold shall ye have : but, behold, thou art a foolish bird ; and when the evil days shall come upon thee, and thou seest the folly of thy heart, return here to me, and I will give thee help." So the king of the hoopoes left the presence of King Solomon with a golden crown upon his head, and all the hoopoes had golden crowns ; and they were exceedingly proud and haughty. Moreover, they went down by the lakes and the pools, and walked by the margin of the water, that they might admire themselves, as it were in a glass. And the queen of the hoopoes gave herself airs, and sat upon a twig ; and she refused to speak to the merops, her cousins, and the other birds who had been her friends, because they were but vulgar birds, and she wore a crown of gold upon her head.

Now there was a certain fowler who set traps for birds ; and he put a piece of a broken mirror into his trap, and a hoopoe that went in to admire itself was caught. And the fowler looked at it, and saw the shining crown upon its head ; so he wrung off its head, and took the crown to Issachar, the son of Jacob, the worker in metal, and he asked him what it was. So Issachar, the son of Jacob, said, " It is a crown of brass," and he gave the fowler a quarter of a shekel for it, and desired him, if he found any more, to bring them to him, and to tell no man thereof. So the fowler caught some more hoopoes, and sold their crowns to Issachar, the son of Jacob ;



until one day he met another man who was a jeweller, and he showed him several of the hoopoes' crowns. Whereupon the jeweller told him that they were of pure gold, and he gave the fowler a talent of gold for four of them.

Now when the value of these crowns was known, the fame of them got abroad, and in all the land of Israel was heard the twang of bows and the whirling of slings; bird-lime was made in every town, and the price of traps rose in the market, so that the fortunes of the trapmakers increased. Not a hoopoe could show its head but it was slain or taken captive, and the days of the hoopoes were numbered. Then their minds were filled with sorrow and dismay, and before long few were left to bewail their cruel destiny.

At last, flying by stealth through the most unfrequented places, the unhappy king of the hoopoes went to the court of King Solomon, and stood again before the steps of the golden throne, and with tears and groans related the misfortunes which had happened to his race.

So King Solomon looked kindly upon the king of the hoopoes and said unto him, "Behold, did I not warn thee of thy folly, in desiring to have crowns of gold? Vanity and pride have been thy ruin. But now, that a memorial may remain of the service which thou didst render unto me, your crowns of gold shall be changed into crowns of feathers, that ye may walk unharmed upon the earth." Now, when the fowlers saw that the hoopoes no longer wore crowns of gold upon their heads, they ceased from the persecution of their race; and from that time forth the family of the hoopoes have flourished and increased, and have continued in peace even to the present day.



THE INDIAN LONG-TAILED NIGHTJAR
(*Caprimulgus macrurus albonotatus*)



THE INDIAN LONG-TAILED NIGHTJAR

(*Caprimulgus macrurus albonotatus*).

Lone on a fir branch his rattle note unvaried
Brooding o'er the gloom, spins the brown eve-jar.

Meredith—*Love in a Valley*.

THE Nightjars form a small family of birds of which eleven species, representing fourteen named forms contained in two genera, occur within Indian limits. One species, the Great-eared Nightjar (*Lyncornis cerviniceps*), occurs in Burma and Travancore and is readily distinguishable by the presence of tufts of feathers just above and behind the ear-coverts, but all our other Indian Nightjars belong to the typical genus *Caprimulgus*, without ear-tufts, and are very much alike, all being brownish or yellowish-grey mottled with darker and lighter spots. In habits, as in general appearance, they are all very similar, resting during the day-time on the ground amongst vegetation or stones or clods of earth, and appearing on the wing only at dusk, when they hawk about for insects which they catch either on the wing by means of their large mouth, which can be opened very widely, or on the ground. They are rarely seen in the day-time, when they keep quite motionless, squatted on the ground, and are readily passed over as a clod of earth, but are commonly seen on the wing in the evening or may be found on roads catching dung-beetles attracted to cattle-droppings.




As in some other groups of birds, the typical Nightjars have a comb-like formation on the inner side of the long middle-toe, but the use of this structure appears to be unknown although it is supposed to be of use in cleaning the beak and the long rectal bristles from the hooked claws of beetles and from the scales of moths which have been caught in the mouth. These birds have also strongly developed rectal bristles which doubtless assist them to obtain insect prey on the wing.

Their noiseless flight is very characteristic and more resembles that of a gigantic moth than that of a bird. It consists of a few quick flappings of the wings alternated with rapid and complicated glides and wheels through the air with wings widely extended. Sometimes the tips of the long wings are brought together above the back with an audible smack. Its position at rest also makes it easy to place a bird as a Nightjar, as it squats down with its whole body on the resting surface; also when a Nightjar perches on a branch, it sits on it lengthwise, and not across it as other birds do.

Their food consists entirely of insects, largely moths and beetles, captured both on the wing and on the ground. They may therefore be considered as useful birds but hardly occur in sufficient numbers, as a rule, to do very much good. It should be noted, however, that the quantity of moths captured and devoured by a single bird is astonishingly large.

As already noted, all the Nightjars look very much alike, at least when seen at large after night-fall. The Common Indian Nightjar (*Caprimulgus asiaticus*), distinguished by having distinct

but narrow black streaks on the back, in combination with an almost naked tarsus, occurs commonly throughout most parts of the Plains of India, Ceylon and North Burma. It is found in open and cultivated country, groves, gardens and non-forested areas generally and is often seen near habitations. It occurs commonly



Common Indian Nightjar (*Caprimulgus asiaticus*); Left foot and head.
(After Blanford, *F. I. Birds*, Vol. III, figs. 53 and 54.)

in the larger gardens of Calcutta and is sometimes known as the "Ice-bird," because its cry, which is constantly heard at night, is like the sound made by a stone skimming over the frozen surface of a pond, the note being repeated slowly at first and then more quickly.

In North Bihar the Common Nightjar seems to be replaced by the Indian Long-tailed Nightjar (*Caprimulgus macrurus albonotatus*), shown on our Plate, which is very similar to the Common Nightjar but rather larger in size, and may be distinguished by having the tarsus feathered throughout. It is found throughout the Plains from North-West India to Assam, extending into the lower ranges of the Himalaya, wherever suitably wooded areas



occur, but in the summer months it ascends higher up the Hills, having been noted at Bhimtal and Masuri.

Regarding its habits, Stuart-Baker, writing on the Birds of North Cachar, gives the following interesting note, which may be quoted *in extenso* :—"My house is built on a hill, the garden on the steepest side coming up to the very brow of the steep, almost precipitous grass slope, leaving room only for a narrow foot-path for the servants and hill-people just outside the fence. This pathway is the favourite haunt after dusk of this fine Nightjar and I, seated motionless on the bank, often have had them approach me within a few feet, so near indeed that I have more than once tried to catch them with a short butterfly-net. I believe it is not at all ageerally known how much these birds feed on the ground, but I have constantly observed them so feeding, and butterflies or other large *dead* insects which were placed near their favourite resting-places were greedily eaten by them. Their movements on the ground are stronger and quicker than might be expected, judging from the formation of their feet, and they *run* in exactly the same manner as do martins and swallows when collecting mud for their nests.

"A very remarkable trait in this bird is the fact that the female will accept the advances of more than one male, but, remarkable as it is, it is undoubtedly the fact, for on one occasion I was a witness of it nor could there be any mistake, for both males were present at the same time and within three yards of me.


"The actions of the young are very peculiar; tiny mites, still blind—their eyes do not seem to open till the seventh day—will,



when first discovered or when they hear a heavy tread near them, lie flat on the ground, their colour closely assimilating with the dead bamboo leaves or other material on which they lie; should, however, the danger of discovery become very imminent, they will crawl under the leaves and hide from sight altogether."

Like other Nightjars, the Indian Long-tailed Nightjar makes no nest, laying its eggs on the bare ground, usually in some sheltered situation. Pairing begins in March and eggs are usually laid by the end of this month or early in April in the Plains, or a little later in the Hills. The number of eggs laid seems to be always two. The eggs vary somewhat in colour, from creamy-white to salmon or fleshy clay-colour, blotched and speckled with reddish-brown, and measure about 30 by 22 mm. The egg is shown in figure 26 of Plate Two at the end of this book. The hen bird sits very closely on her eggs, so that she may almost be trodden upon before flying off. Speaking of Nightjars in general, Newton remarks:—"So light is it that the act of brooding, even where there is some vegetable growth, produces no visible depression of the grass, moss, or lichens on which the eggs rest, and the finest sand almost equally fails to exhibit a trace of the parental act. Yet scarcely any bird shows greater local attachment and the precise site chosen one year is almost certain to be occupied the next."

The call of this species is quite different from that of the "Ice-bird," consisting of the sounds "*chouunk chouunk*" repeated at intervals, and may be compared to the sound made by striking a plank with a hammer. When on the wing it also utters a low chirping cry.





THE HAWK-CUCKOO
(*Hierococcyx varius*)



THE HAWK-CUCKOO

(*Hierococcyx varius*).

The merry cuckoo, messenger of Spring,
His trumpet shrill hath thrice already sounded.

Spenser.

And higher in the uplands we may hear,
Greeting the Spring through scented pinewood trees,
Faint echoes sweet—the cuckoo calling clear
Mingling with murmurs of the mountain bees.

Chenar Leaves.

THE common name "Hawk-Cuckoo" conveys a good description of this bird, as it is really a Cuckoo which looks very like a hawk. It is about the size of a myna, but with a longer tail, greyish-brown in colour, whitish beneath, the breast tinged with pink, each feather with darker cross-bars, eyes and legs brilliant yellow. When on the wing, it looks very much like a small hawk but, when it alights, it at once assumes a slouching, cuckoo-like attitude, with the wings dropped forward so as to touch the perch and the tail slightly raised and expanded, thus presenting an aspect very different from the compact and alert look of a hawk. Seen thus, at rest, this bird can hardly be mistaken for a true hawk, as it has the furtive, peering ways of common cuckoos, constantly jerking itself from side to side and puffing out its throat.

The appearance of the Hawk-Cuckoo is probably less familiar to most people than is its note, which has aptly earned for it the notorious title of the "Brain-Fever Bird." Our Indian gardens and groves contain many sweet-voiced singers amongst their avian denizens and a few whose voices are less grateful to the ear, but

there is not one whose notes consist of such ear-splitting and nerve-racking cries as do those of the Brain-Fever Bird. With the most annoying persistence and reiteration this bird repeats its cry, which bears a remarkable resemblance to the word "brain-fever" repeated in a piercing shriek running up the scale. The cry may also be



Head and foot of Hawk-Cuckoo (*Hierococcyx varius*).

written as "Pipiha" and in some districts the vernacular name of the bird is given as *Pupiya*. Another rendering of the call, which includes the overture preceding the triple note, is, "O lor'! O lor'! how very hot it's getting—we feel it, *we feel it*, WE FEEL IT."



The call is extremely loud and shrill and can be heard—indeed, it cannot but be heard—within a radius of several hundred yards, but one of the most annoying things about it is its intermittent character. The human ear soon becomes accustomed to any continuous and uniform kind of noise. One becomes so accustomed to the buzz of a dynamo that one awakens at once if it stops. The copper-smith *tonk-tonking* in the garden all day is hardly heard consciously unless one listens for it. But the shrieks of the Brain-Fever Bird burst their way without ceremony into one's inner consciousness, whether awake or asleep, and one cannot help but hear them. “We feel it, *we feel it*, WE FEEL IT” go the cries, up and up the scale, and then suddenly stop, and one hopes fervently that this fiend in bird's plumage has burst its throat or at least flown away out of ear-shot. But no; after a short interval it begins again and may continue for hours at a stretch. Very often the performance commences just at dusk, when it has got too dark to make out the culprit, and lasts all night without intermission. When this sort of thing takes place on a really hot night, the victim, who is attempting to woo sleep after a hard day's work, may well be excused if the first dim dawn sees him sallying forth on vengeance bent. But vengeance is not always easy to attain. The bird usually perches high up in a tall tree and keeps so still and is so inconspicuously coloured that, even when its shrieks locate the very branch whereon it is sitting, it is not always easy to make out. Furthermore, it is wary and often flies off as soon as it sees that it has been detected. There are, however, usually only a few individuals in each locality and a comparatively small reduction in numbers works wonders in abating the nuisance. The call being very penetrating, it often happens that these birds call to one another across a distance of perhaps half a mile and, by shooting one bird forming a link in the chain between others on either side of it, the chain is broken, and a blessed peace reigns once more, at least until another bird invades the immediate neighbourhood. In Bihar the call of this bird coincides with the approach and duration of the hot, dry



weather before the monsoon ; occasionally it may be heard as early as in December but more usually commences about February and is continued, becoming more frequent and continuous, until the Rains break, when there is a welcome cessation for a few months. In other districts this may not be so ; thus, as regards Calcutta, Cunningham states that " there is hardly any season at which their characteristic notes may not occasionally be heard ; but, as a rule, it is during the rainy months that they are most frequent, so that the designation ' hot-weather bird,' that is often applied to the species in other parts of the country, is hardly applicable to it in Calcutta."

According to Blanford, the Hawk-Cuckoo occurs throughout the whole of India and Ceylon, extending as far East as Dacca, but not to Assam, and West to Rajputana, but not to Sind or the Punjab ; but, although odd examples may occur throughout this area, its range as a common bird seems to be more restricted. It is extremely common in the United Provinces and Bihar, less common further south in Bengal. Dewar notes that he never heard it in Madras, nor did I ever hear it during my residence in Coimbatore, and it is apparently quite absent in the island of Bombay. In some districts in which it is absent, or at least scarce, the Hawk-Cuckoo is frequently confounded with the Koël and the name " Brain-Fever Bird " given to the latter. As Dewar puts it, " There is certainly some excuse for the mistake, for both are cuckoos and both are exceedingly noisy creatures ; but the cry of the koël bears to that of the brain-fever bird or hawk-cuckoo much the same relation as the melody of the organ-grinder does to that of a full German band. Most men are willing to offer either the solitary Italian or the Teutonic gang a penny to go into the next street, but, if forced to choose between them, select the organ-grinder as the lesser of the two evils. In the same way, most people find the fluty note of the koël less obnoxious than the shriek of the hawk-cuckoo."



In spite of its obnoxious vocal efforts, the Hawk-Cuckoo does some little good by feeding on injurious insects, although when it can find time to hunt these out in the height of the hot weather, when it seems to be calling continuously day and night, always seems somewhat of a mystery. Like other cuckoos, it eats hairy caterpillars, whose defensive armament protects them from the attacks of most other birds, and it also eats other caterpillars, crickets, grasshoppers, bugs and beetles. The diet is a mixed one, comprising buds and fruits, particularly wild fig fruits, as well as insects. It is presumably on account of this redeeming feature that this bird is protected in Delhi, the United Provinces, Bengal and Assam (where, however, it does not occur!).

Like other Cuckoos, the Brain-Fever Bird economizes in house-keeping, building no nest of its own but placing its eggs in the nests of other birds, usually the "Seven Sisters" or some allied species of Babblers, the breeding-season lasting from April until June. The eggs are deep blue in colour and measure about 26 mm. by 20 mm., and are about the same size and shape as those of the foster-parents. The Babbler's eggs are wholly blue, very glossy, hard-shelled and broad, blunt ovals in shape; the Hawk-Cuckoo's eggs are very similar in colour but with a softer, more satiny surface, less glossy and with much thicker shells, in shape rather more spherical or elliptical and slightly larger than in the Babblers. When lying side by side in the nest, however, the eggs of the Babbler and of the parasitic Cuckoo are often practically indistinguishable.

The manner in which Cuckoos' eggs are deposited in the nests of other birds is one which has engaged a great deal of attention. It used to be supposed that the eggs were laid in the normal way in the nest of the birds selected as foster-parents and this may occasionally be done, but the more frequent method is for the egg to be laid and then carried by the cuckoo in its bill and dropped into the nest which has been selected for the purpose. The unusually

thick texture of the cuckoo egg-shell seems to be specially adapted to this end as, in cases where the nest is placed inside a hole, the egg may have to be dropped into it from a little height. In the case of the Hawk-Cuckoo, it is possible that its hawk-like appearance on the wing may be advantageous in securing a clear field for depositing an egg in this way in the nest of the Seven Sisters, as one observer states that the whole sisterhood make themselves scarce when the Hawk-Cuckoo appears on the scene, and thus give her a fair field for planting her oval imposition on them. Our Plate shows a Hawk-Cuckoo, with an egg in her bill, about to be dropped into the nest of one of these Babblers. It may be added that further observations, on the method of egg-deposition employed by this and other Indian Cuckoos, are very desirable.

The Hawk-Cuckoo is known vernacularly as *Kupak* and *Pupiya* in Hindi-speaking districts, as *Chok-gallo* in Bengal, as *Zakhat* in the Deccan, as *Irolan* in Malayalam and as *Kutti-pitta* in Telugu districts.



Spotted Dove.



THE ROSE-RINGED PAROQUET

(*Psittacula torquata*).

.. . . . Bold parrots hither wing,
And settle down, with scream and chattering.

Nebula to Man.

The plum-necked parrots swung from fruit to fruit.

The Light of Asia.

Ye, too, who dwell so merry in the trees—,
Quick-darting parrots, bee-birds, bulbuls, doves—,
None of ye hate your life, none of ye deem,
To strain to better by foregoing needs.

The Light of Asia.

Of wild fruit fallen from the boughs o'er head,
Shaken to earth by chattering ape or plucked
By purple paroquet.....

The Light of Asia.

THE great family of the Parrots is so widely distributed throughout the Tropical Regions of the World and is composed of so many different species that it is difficult to say why the Indian list of indigenous forms should be so scanty, only a dozen species, represented by nineteen sub-species, being known to occur. But perhaps it is well that we have no more; for, as Dewar remarks, "the green parrot is one of those good things of which it is possible to have too much."

Parrots are sharply distinguished from all other classes of Birds by several anatomical peculiarities in their vertebræ, feet, and other parts, into which we need not enter here. Most Indian Parrots are easily recognizable as such, their most obvious characters being the short, stout, strongly-hooked bill, thick fleshy tongue, movable upper mandible, climbing habits, and (in most species)

long tail and greenish colour. The subject of our present article may be distinguished from all other Indian Parrots by its long tail, head (except chin and mandibular stripe in males) and body green, bill deep red, with no red patch on the wing-coverts. It is by far the commonest and most familiar of the green Indian Parrots, occurring abundantly in all open and cultivated land around towns and villages, often in large flocks, dashing over the tree-tops in swift, arrowlike flight. It is found practically everywhere throughout the Plains of India, Burma and Ceylon, but seems to be commoner in Northern India than in Madras.



Rose-ringed Paroquets attacking wild fruits.

Wherever it occurs, however, in its wild state the Rose-ringed Paroquet is an unmitigated nuisance, as its diet is wholly vegetarian and it feeds largely on cultivated grains and fruits. When a large flock descends on a ripening crop of *guar* (sorghum) or similar cereal,

THE ROSE-RINGED PAROQUET

151

a great deal of damage is done, some by the actual grain that is eaten, but far more by the extremely wasteful method of feeding of this bird, which often breaks off a whole head, delicately



Sugarcane damaged by parrots in Assam (From a photograph by Dr. C. A. Barber).

selects one or two grains, throws away the rest, and breaks off another head which is treated in the same way. When fruits are ripe, these birds soon find them out and play havoc with them.



When no cultivated fruits or crops are in season, the food consists of wild fruits (wild figs, *Zizyphus*, etc.) and seeds. The late Mr. C. W. Mason examined fifty-three birds at Pusa and Mr. D'Abreu three more at Nagpur, and in all cases the stomach-contents consisted entirely of vegetable matter—mustard, wheat, maize, paddy, litchi and wild fruits, and seeds of *Dalbergia sissu*. When the silk-cotton trees are in flower in February these parrots are amongst the crowd of birds which congregate to imbibe the nectar. We have not yet had any complaints of its attacking sugarcane but, with the increasing cultivation of this crop in Bihar, it will perhaps discover that it is edible and attack it in the same way as another Parrot has damaged sugarcane in Assam, by gnawing large holes in the stems.

The Rose-ringed Paroquet does not deserve, nor has it been afforded, any protection under the Wild Animals Protection Act: on the contrary, its destruction should be encouraged and, if this could be done by exempting the export of its plumage from the present Customs restrictions, or in any other way, it would be all to the benefit of the cultivator who is practically powerless at present to prevent the immense damage done to his crops by this destructive bird.

It must be admitted, however, that from a purely æsthetic and non-utilitarian view-point the Rose-ringed Paroquet is a decidedly handsome and attractive bird when seen under natural conditions flashing through the sunshine or climbing over a tree in search of food, and, in places where its numbers are but small, its brilliant hues, delicate outlines and dainty ways provide a perennial source of joy to the bird lover. But in country districts, where it is only too abundant, its destructive habits and shrill harsh screams very quickly nullify such feelings of admiration.

As a cage-bird the Rose-ringed Paroquet is a favourite all over India, and thousands of young birds must be taken every year



from their nests and doomed to pass the remainder of their existence within prison bars, although unfortunately this annual toll seems to make little difference to the total numbers which exist to ravage the farmer's crops. As its powerful beak soon secures its escape from a wooden cage, this bird is usually kept in a small dome-shaped cage made of hoop-iron, with a sheet-iron floor, which must get uncomfortably like an oven in the hot weather. With a little instruction, this parrot often becomes a good talker, and it can also be taught to perform tricks. Lockwood Kipling tells of one that he used to see in the streets of Delhi "that went through gymnastic and military exercises, whirling a tiny torch lighted at each end, loading and firing a small cannon, lying dead and coming to life again; all done with a comic air of eagerness and enjoyment which it seemed hard to impute to mere hunger for the morsels that rewarded each trick."

The Rose-ringed Paroquet breeds in January and February in Southern India, from March to May further North. No regular nest is made but the eggs are laid in a branch of a tree, occasionally in a wall. If a suitable hole is already available, the birds appropriate it, and couples may often be seen, at Pusa about the end of February every year, inspecting eligible sites, which by that time are often occupied by Hoopoes. If no acceptable hole can be found, the birds may excavate one for themselves and Colonel Butler noted a pair at Deesa which were at work clearing out the hole, in which the eggs were subsequently deposited, for at least three months before the eggs were laid. A finished hole, either acquired or excavated either wholly or in part, is generally about two inches in diameter, and goes straight into the trunk for two to four inches and then turns downward for a distance of six inches to three feet, the lower portion being expanded into an egg-chamber which is four or five inches in diameter. Should a natural hollow in the tree be utilized or cut into, the chamber may be much larger. The entrance-hole is cut either into the trunk of a tree or into a

large bough ; in the latter case it is often placed on the lower side of the bough. No lining is provided, the eggs being laid on a few chips of wood at the bottom of the hole. The usual number of eggs is four but as many as six are found at times. The egg is pure white, without any gloss, usually in shape a moderately broad oval, considerably pointed towards one end, and measures about 30 by 24 millimetres.

Our Plate shows both sexes of this Paroquet, the adult male being distinguishable by his rosy collar which is absent in the female.



Head of male Rose-ringed Paroquet.



THE INDIAN SPOTTED OWLET
(*Carine noctua brama*)



THE INDIAN SPOTTED OWLET

(*Carine noctua brama*).

... Disquiet yourself not;
'Tis nothing but a little downy owl.

Shelley.

... Here owls waylay
Snails, beetles, rats, and other passing prey.

Nebula to Man.

Over the shining pavements peacocks draw
The splendours of their trains, sedately watched
By milk-white herons and the small house-owls.

The Light of Asia.

OF the two score or so of species of owls which are found within Indian limits, the Indian Spotted Owlet (*Carine noctua brama*) is probably at once the commonest and most familiar, as it occurs abundantly throughout the Plains of India North of 14° latitude and Northern Burma and is in evidence during the early evening often before sunset and long before dusk, when most other owls have not yet ventured out. It is, moreover, quite a domestic owl, keeping especially to trees in cultivated tracts such as gardens around houses, and it is commonly found roosting and breeding in the roofs of houses where these afford the necessary shelter. It is fond of perching on the branch of a tree or on a pole or fence or telegraph-wire; indeed, as Hume remarks, it is one of the birds that seem to think that telegraph-wires were erected for their sole and especial benefit. It is a decidedly noisy bird, making itself most evident to the ear in the evening or early morning, but frequently heard at intervals during the night, when a regular volley of chuckles and squeaks is poured forth by one or more of these little owlets; but the noise is not sustained for any length of time.

In the day-time, like other owls, the Spotted Owlet hides away in some dark corner, such as a hole in a tree or wall or in a house-roof



or even in a bungalow verandah if no better situation offers, emerging towards sunset to hunt for prey. As Cunningham remarks, it is most diverting to watch them emerging; one after another, before fairly coming out, putting forth its queer little round head and staring eyes through the opening of the cavern. After they have emerged they usually sit very quietly for a time as though only half awake, and are either perfectly silent or occasionally utter a low-toned "chirrk." Then, all of a sudden, they begin to chuckle and finally break out into a perfect torrent of hoarse chattering; and finally, after having indulged in such exercises for some minutes, they spread their short, rounded wings and sail off to their night's hunting.

However, it does not seem to mind the full blaze and heat of the sun, as we have seen it in the early afternoon on a hot April day sunning itself with outspread wings on the bare gravel in front of the bungalow. A pair which have for years inhabited the bungalow of one of us (C. M. I.) may often be seen during the day-time either perched on some *sambhar's* antlers on the verandah wall or else flying with undulating flight to a tree in the garden, usually to a tamarind tree, and, after staying there a short time, returning to their hole. We have also seen them on a drizzly day seated on a perch outside, enjoying a shower bath.

It is one of the birds that hawk termites (white ants, so called) along with Rollers, etc. Mr. Stuart-Baker writes that they are great bat hunters, not catching them on the wing but hauling them out of their holes; but apparently all these owls do not wage war against bats and this habit is perhaps local or confined to a few individuals.

It lives chiefly on insects and to a less extent on mice, shrews and lizards, its insect prey consisting mostly of beetles and crickets. In the case of eight birds examined at Pusa by the late C. W. Mason, of 69 insects taken, one was of a beneficial species, twenty-six were



neutral and forty-two were injurious. Hume mentions a case, reported by Colonel Butler, in which a pair of Spotted Owlets had apparently attacked a nesting Paroquet (*P. torquata*), killed it on the nest and taken possession of this for themselves. As the Paroquet is a perfect pest to the agriculturist, we can only wish that the Spotted Owlet would act regularly in this way. At Pusa this little owl seems to live largely on large crickets (*Brachytrupes* and *Gryllotalpa*) and on dung-beetles. There is no doubt that it is a most useful bird in all districts where large crickets occur so commonly as to do damage. As these crickets are nocturnal, retreating into their subterranean burrows during the day-time and only emerging to feed under cover of night, their most effective natural enemies are those, such as the Spotted Owlet, which are also nocturnal and which can swoop down and destroy them whilst they are feeding above-ground. One commonly sees this little owl swoop down from its perch and catch some prey on the ground, returning to its perch to devour it, usually to the accompaniment of a burst of chattering which is presumably its form of thank-offering for a good supper.

Our Plate gives a good idea of this little owl and the left-hand figure shows the terrifying attitude, assumed, after first sitting up very erect, by suddenly crouching down and frowning and glaring in a terrible way, to frighten any observer or intruder.


The Spotted Owlet breeds from February to April, the period being slightly earlier in the South and later in Northern India, but the great majority of birds lay in March, laying usually three, four or five white (pink when fresh) eggs in a hole, in a tree or building, the nest being scantily lined with a few dry leaves, grass, decayed wood, or feathers. We have taken three clutches of three eggs each from one nest but even then the birds would not desert their nesting site. Incubation evidently starts as soon as the bird lays as we have taken at the same time from one nest one young, two highly incubated and one perfectly fresh egg. The fluffy young,




when fledged, are drab-and-white with yellow eyes just like their parents and, also like their parents, are exceedingly noisy, each member of the family, as Dewar puts it, talking gibberish at the top of its voice, sixteen to the dozen, and as all will persist in speaking at once, the result is a nocturnal chorus that will bear comparison with the efforts of the cats which enliven the Londoner's back yard.

Jerdon, quoting Sykes, writes that "the Mahrattas have a superstition regarding this species, and a class of people called *Pingleh* live on the credibility of the people by pretending to consult it and predict events." Jerdon also says that this Owlet is used by some *shikaris* to catch small birds. It is pegged out near a low bush, the branches of which have been smeared with bird-lime, so that any small birds that come to mob it settle on the bush and are caught on the limed twigs. It is, however, noticeable that this owl does not seem to be molested by birds, as other larger owls are, probably because it is more at home in daylight and so does not attract notice by the blundering flight assumed by other owls when they venture out by day. Its flight is indeed far from a blundering one and, in a recent issue of the Bombay Natural History Society's Journal, Mr. G. O. Allen has called attention to its occasional habit of hovering; this we have also noticed at times and it is probably assumed when watching some small prey which has emerged from its burrow.


Besides the North Indian form five other geographical races of this little owl occur within Indian limits; these are the Southern Spotted Owlet (*C. noctua fryi*), found in Southern India below 14° North latitude; the Burmese Spotted Owlet (*C. noctua pulchra*), found in Central and Southern Burma; the Forest Spotted Owlet (*C. noctua blewitti*), found in forest areas in Sambalpur, Karial and Khandesh; Hutton's Owlet (*C. noctua bactriana*), a straggler into North-West India along the Afghan frontier; and the Baluchistan Spotted Owlet (*C. noctua tarayensis*), found in Sind and Baluchistan.




1




2




3




4



5




6




7




8




9




10




11




12



13




14



15




16




17




18




19




20




21




22




23




24




25




26




27



28



29



30

PLATE II.

- Fig. 19. GOLDEN-BACKED WOODPECKER White, smooth, slightly glossy.
 „ 20. GREEN BARBET Dull white, smooth, not glossy.
 „ 21. COPPERSMITH Pure white, smooth.
 „ 22. INDIAN ROLLER Pure white, glossy.
 „ 23. COMMON INDIAN BEE-EATER.. Pure white, glossy.
 „ 24. BLUE-TAILED BEE-EATER .. Pure white, glossy.
 „ 25. INDIAN HOOPOE Dull dirty white, no gloss.
 „ 26. INDIAN LONG-TAILED NIGHTJAR Glossy, whitish slightly tinged with
 yellow, with pale blotches of brown
 and purple.
 „ 27. HAWK-CUCKOO Smooth, dull blue.
 „ 28. ROSE-RINGED PAROQUET .. Dull white, smooth, without pitting.
 „ 29. INDIAN SPOTTED OWLET .. Dead white, smooth, hardly glossy.
 „ 30. CATTLE EGRET Dull white, smooth, not glossy.

Note. The short descriptions of Eggs, given above, are taken from the actual (blown) specimens from which the drawings were made.

865